

Policy Number	RX501.076
Policy Effective Date	01/01/2026
Policy End Date	07/31/2026

Testosterone Replacement Therapies

Table of Contents
Coverage
Policy Guidelines
Description
Rationale
Coding
References
Policy History

Related Policies (if applicable)
SUR717.001 Gender Assignment Surgery and Gender Reassignment Surgery and Related Services
RX501.007 Hormone Replacement Therapies (HRT) Using Implanted Pellets for Women and Delayed Puberty

Disclaimer

Medical policies are a set of written guidelines that support current standards of practice. They are based on current generally accepted standards of and developed by nonprofit professional association(s) for the relevant clinical specialty, third-party entities that develop treatment criteria, or other federal or state governmental agencies. A requested therapy must be proven effective for the relevant diagnosis or procedure. For drug therapy, the proposed dose, frequency and duration of therapy must be consistent with recommendations in at least one authoritative source. This medical policy is supported by FDA-approved labeling and/or nationally recognized authoritative references to major drug compendia, peer reviewed scientific literature and generally accepted standards of medical care. These references include, but are not limited to: MCG care guidelines, DrugDex (IIa level of evidence or higher), NCCN Guidelines (IIb level of evidence or higher), NCCN Compendia (IIb level of evidence or higher), professional society guidelines, and CMS coverage policy.

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Legislative Mandates

EXCEPTION: For members residing in the state of Ohio, § 3923.60 requires any group or individual policy (Small, Mid-Market, Large Groups, Municipalities/Counties/Schools, State Employees, Fully-Insured, PPO, HMO, POS, EPO) that covers prescription drugs to provide for the coverage of any drug approved by the U. S. Food and Drug Administration (FDA) when it is prescribed for a use recognized as safe and effective for the treatment of a given indication in one or more of the standard medical reference compendia adopted by the United States Department of Health and Human Services or in medical literature even if the FDA has not approved the drug for that indication. Medical literature support is only satisfied when safety and efficacy has been confirmed in two articles from major peer-reviewed professional medical journals that present data supporting the proposed off-label use or uses as generally safe and effective. Examples of accepted journals include, but are not limited to, Journal of

American Medical Association (JAMA), New England Journal of Medicine (NEJM), and Lancet. Accepted study designs may include, but are not limited to, randomized, double blind, placebo controlled clinical trials. Evidence limited to case studies or case series is not sufficient to meet the standard of this criterion. Coverage is never required where the FDA has recognized a use to be contraindicated, and coverage is not required for non-formulary drugs.

Coverage

This medical policy has become inactive as of the end date above and is not to be used for current claims adjudication or business purposes. Testosterone replacement therapy is now addressed in medical policy RX501.007.

This medical policy does NOT address Gender Reassignment Services (Transgender Services). This medical policy IS NOT TO BE USED for Gender Reassignment Services. Refer to SUR717.001, Gender Assignment Surgery and Gender Reassignment Surgery with Related Services.

Testosterone replacement therapy **may be considered medically** under the following conditions:

- An established diagnosis of hypogonadism with androgen deficiency (see Policy Guidelines section) that includes:
 - Persistently low total testosterone levels; **and**
 - Multiple symptoms of hypogonadism including at least 1 “more specific” symptom (see Policy Guidelines section).
- Human immunodeficiency virus-infected men (biological sex) with low testosterone levels and weight loss; **OR**
- Men (biological sex) on chronic steroid treatment with low testosterone levels (see Policy Guidelines section).

EXCEPTION: Individual consideration may be given for men continuously on therapy for at least one year and a morning total serum level or free testosterone level tested within the last year remains below or within the testing laboratory’s normal range.

Testosterone replacement therapy **is considered experimental, investigational and/or unproven** in all other situations in which the above criteria are not met, including but not limited to older men (biological sex) with type 2 diabetes mellitus and androgen deficiency or low testosterone levels in the absence of clinical signs and symptoms of hypogonadism.

Policy Guidelines

Diagnosis of Androgen Deficiency

An established diagnosis of hypogonadism with androgen deficiency includes appropriate evaluation and diagnostic workup of a man who presents with symptoms of hypogonadism. Screening in asymptomatic populations is not recommended. Measurement of serum total testosterone is initially used; serum-free testosterone levels can be measured when total

testosterone is in the low to normal range, and alterations of serum hormone-binding globulin are suspected. (1) Once a persistently low testosterone level has been established, diagnostic testing of the hypothalamic-pituitary axis should be performed to distinguish primary hypogonadism from secondary hypogonadism. When secondary hypogonadism is identified, the underlying etiology should be identified, and any reversible causes treated appropriately before consideration of testosterone replacement.

Men on chronic steroid treatment would be receiving ongoing treatment for a chronic condition as opposed to episodic treatment for an acute condition or acute flare of a chronic condition. The length of acute episodic steroid treatment may vary from several days to several months, but in most cases, will be less than 4 to 6 weeks.

Persistently low testosterone levels refer to serum levels below the lower limit of normal on at least 2 occasions when measured in the early morning. The threshold lower limit for serum testosterone levels is not standardized. The Endocrine Society has recommended a lower limit for normal levels of 300 ng/dL for total testosterone and 9.0 ng/dL for free testosterone. (1) Joint guidelines from several European and American specialty societies have recommended that replacement therapy be considered at serum total testosterone levels less than 350 ng/dL.

“More specific” symptoms of hypogonadism, as classified by the Endocrine Society, include the following (1):

- Incomplete or delayed sexual development
- Decreased libido
- Decreased spontaneous erections
- Breast discomfort, gynecomastia
- Loss of axillar and/or pubic body hair
- Very small (<5 mL) or shrinking testes
- Infertility due to low sperm count
- Height loss due to vertebral fractures, low trauma fractures, low bone density
- Hot flushes, sweats.

Monitoring Strategies for Individuals on Testosterone Therapy

Monitoring of testosterone replacement should be performed beginning 3 to 6 months after replacement is initiated to ascertain whether serum levels are restored to the normal range, to determine whether clinical symptoms have improved, and to monitor for adverse events. The goal of testosterone replacement is to raise levels into the mid-normal range. Higher replacement levels are unlikely to improve symptoms further and may increase the incidence and/or severity of adverse events.

Recommendations for monitoring for testosterone-related adverse events have been provided by the Endocrine Society guidelines on testosterone therapy in men with androgen deficiency (see Practice Guidelines and Position Statements section). (1) These recommendations include:

- Determine hematocrit levels at baseline, at 3 to 6 months, and then annually. If the hematocrit level is above 54%, stop therapy until the hematocrit level decreases to a safe level, evaluate the individual for hypoxia and sleep apnea, and reinstitute therapy at a reduced dose.
- Repeating bone mineral density of the lumbar spine, femoral neck, and hip after 1 to 2 years of testosterone therapy in hypogonadal men with osteoporosis.
- For men 55 to 69 years of age, and for men 40 to 69 years of age who are at increased risk for prostate cancer, conduct a digital examination of the prostate and prostate-specific antigen (PSA) measurement before initiating treatment, at 2 to 3 months after initiating treatment, and then in accordance with evidence-based guidelines for prostate cancer screening, depending on the age and race of the individual.
- Obtain urological consultation if there is:
 - An increase in serum or plasma PSA concentration greater than 1.4 ng/mL within 12-months of initiating testosterone treatment.
 - A confirmed PSA of more than 4 ng/mL at any time.
 - Detection of a prostatic abnormality on digital rectal examination.
 - Substantial worsening of lower urinary tract symptoms.

Description

Testosterone replacement therapy is the primary treatment for androgen deficiency in men. Testosterone replacement is intended to counter the adverse effects of low testosterone levels with clinical signs and symptoms of hypogonadism. A variety of testosterone preparations are available for clinical use.

Testosterone and Testosterone Levels

Testosterone is produced in males, primarily by the testes, in response to stimuli from the hypothalamic and pituitary glands. Low testosterone is caused by deficient production of the hormone and is also known as androgen deficiency. Primary androgen deficiency results from failure of testosterone production at the testicular level in the presence of normal hypothalamic and pituitary function. Secondary androgen deficiency results from failure of the pituitary gland to produce androgen-stimulating hormones (luteinizing hormone, follicle-stimulating hormone). It can be caused by dysfunction at the hypothalamic or pituitary level.

Hypogonadism

Diagnosis

Hypogonadism is the clinical syndrome associated with androgen deficiency. The signs and symptoms of hypogonadism depend on the age of onset. In prepubertal males, the hallmark of androgen deficiency is the failure to develop secondary male sex characteristics. In adults, the signs and symptoms are nonspecific, with the most specific symptoms related to sexual functioning such as decreased libido and erectile dysfunction. Symptoms are dependent on age, the severity of androgen deficiency, duration of androgen deficiency, sensitivity to androgen, and comorbid illness. (2) Symptoms and signs other than sexual dysfunction include loss of

body hair, hot flashes or sweats, decreased energy, depression, sleep disturbance, reduced muscle mass and strength, and/or increased body fat. All can occur in the absence of androgen deficiency and, therefore, the diagnosis of hypogonadism can be challenging. A systematic review by Zarotsky et al. (2014) reported on risk factors, comorbidities, and consequences of male hypogonadism identified multiple comorbid conditions that are consistently risk factors for hypogonadism, including advanced age, obesity, metabolic syndrome, and poor general health status. (3) Multiple other conditions, including diabetes, coronary heart disease, hypertension, stroke, and peripheral artery disease, correlated with the presence of hypogonadism, although these were not identified as risk factors.

Testosterone levels decrease with age, beginning in the fourth or fifth decade of a person's life, and this decrease is sometimes referred to as male "andropause." In the European Male Aging Study of 3220 men, there was a decline in serum testosterone levels of 0.4% per year between the ages of 40 and 70. (4) Because this decline is gradual and modest, the clinical impact is uncertain. While there are also parallel decreases in androgen-dependent factors with age, such as sexual function, lean body mass, and bone mineral density, the degree to which these changes are due to decreasing testosterone has not been determined with certainty.

Because of the decline in testosterone levels with age, more elderly males will have lower levels than younger men. Using a cutoff of 325 ng/dL as the lower limit of normal testosterone levels, Travison et al. (2007) estimated, based on a prospective cohort of 890 men, that the rate of low testosterone is 20% for men in their 60s; 30% for men in their 70s; and 50% for men in their 80s. (5) In this study, other factors were associated with decreased testosterone, such as obesity and severe emotional stress. A much lower percentage of men have a combination of low testosterone levels and definite symptoms of hypogonadism. In the European Male Aging Study, this was estimated to be present in 2.3% of men when using a cutoff of at least 3 symptoms potentially related to androgen deficiency.

Another factor that makes the diagnosis of hypogonadism challenging is the measurement of testosterone levels. Testosterone levels fluctuate substantially due to various factors. There is a diurnal variation, which is more pronounced in younger men, with peak levels occurring in the early morning. This makes the timing of measurement important and requires repeated measurement before making a determination that testosterone is consistently low. Also, there is a wide range of levels seen in healthy men and assigning the proper age-appropriate cutoff is controversial. Some men exhibit clear symptoms of hypogonadism with testosterone levels that are in the low to normal range, while other men with low levels do not experience any symptoms.

Treatment

There are numerous U.S. Food and Drug Administration (FDA)–approved testosterone formulations available for replacement therapy. For most delivery preparations, approval was based on the ability to increase levels to the normal range, not to demonstrate beneficial clinical outcomes. (6)

The following are the different preparations and routes of administration:

- Oral Testosterone: The only commercially available oral testosterone products include testosterone undecanoate. There are currently 3 branded products which are not substitutable with one another. Oral testosterone is readily absorbed from the intestine and is rapidly metabolized by the liver. The rapid metabolism in the liver limits its clinical utility because it is difficult to maintain steady serum levels. In addition, the first pass through the liver may increase the probability of liver toxicity.
- Nasal Gel: Testosterone may be administered as a nasal gel, but this dosage form is limited by frequent administration (3 times daily).
- Intramuscular Testosterone: Testosterone undecanoate is an intramuscular (IM) depot preparation that is slowly absorbed into the circulation. It is administered by deep IM injection every 10 to 14 weeks and thus has the advantage of infrequent dosing. Disadvantages of this preparation include the IM injection route, which can be painful, and also inconsistent rates of absorption. Inconsistent absorption can lead to fluctuating testosterone levels and related clinical symptoms. Other intramuscular testosterone products include testosterone enanthate and testosterone cypionate.
- Topical Patch: Topical testosterone patches can be applied to non-genital skin areas. Patches are generally dosed once per day and result in stable testosterone levels over time. A limiting factor of patch use is the development of skin irritation at the patch site in a high percentage of users.
- Topical Gels: A number of topical testosterone gel preparations are commercially available. They range in strength from 1% to 2% and provide stable serum levels. The gel is applied daily on non-genital skin areas. Precautions need to be taken to avoid transmission of the drug to others by direct contact. Therefore, it is recommended that the gel is placed on covered skin and that hand washing is performed after application.
- Subcutaneous Pellets: Another depot formulation is a subcutaneous testosterone pellet. The pellets are placed subcutaneously in the buttocks, abdominal wall, or thigh under local anesthesia. They are replaced every 3 to 6 months. Limitations include the need for minor surgical procedures, and local reactions at the implantation site (e.g., infections, fibrosis).
- Subcutaneous Solution: Testosterone enanthate and testosterone cypionate can be administered subcutaneously once weekly.

Regulatory Status

Numerous preparations of testosterone have been approved by the FDA for use in testosterone replacement therapy. They include IM, oral, topical, subcutaneous, nasal, and buccal (no longer available in the U.S.) preparations.

In March 2015, the FDA issued a drug safety communication for prescription testosterone products. (7) The communication stated: “We are requiring that the manufacturers of all approved prescription testosterone products change their labeling to clarify the approved uses of these medications. We are also requiring these manufacturers to add information to the labeling about a possible increased risk of heart attacks and strokes in patients taking testosterone. Health care professionals should prescribe testosterone therapy only for men

with low testosterone levels caused by certain medical conditions and confirmed by laboratory tests.” The communication also stated: “FDA has concluded that there is a possible increased cardiovascular risk associated with testosterone use. These studies included aging men treated with testosterone. Some studies reported an increased risk of heart attack, stroke, or death associated with testosterone treatment, while others did not.”

As of February 2025, following the FDA’s evaluation of a large clinical trial (TRAVERSE) that found testosterone therapy does not raise the risk of heart attack or stroke compared to a placebo, the agency is removing this risk from the Boxed Warning on all prescription testosterone products. Additionally, a warning about the potential for increased blood pressure is being added to the prescribing information for testosterone products that currently lack this information, based on findings from separate blood pressure monitoring studies. (8)

Rationale

Medical policies assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function—including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

There is a large body of literature that evaluates the efficacy of testosterone replacement therapy. This body of evidence is primarily characterized by small- to medium-sized trials of short duration. There is a high degree of variability in the patient populations, dosing and delivery methods for testosterone replacement, and outcomes measured. There are also numerous systematic reviews of the available evidence. This policy focuses on the impact of testosterone on specific symptoms for adults with androgen deficiency and clinical symptoms of hypogonadism, and on the benefit for specific subpopulations. The discussion emphasizes the available systematic reviews and larger individual RCTs.

Androgen Deficiency and Clinical Symptoms of Hypogonadism

Clinical Context and Therapy Purpose

The purpose of testosterone replacement therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with androgen deficiency and clinical symptoms of hypogonadism.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with androgen deficiency and clinical symptoms of hypogonadism.

Interventions

The therapy being considered is testosterone replacement therapy.

Testosterone replacement therapy is the primary treatment for androgen deficiency in men. Testosterone replacement is intended to counter the adverse effects of low testosterone levels.

There are numerous U.S. Food and Drug Administration approved testosterone formulations available for replacement therapy.

Comparators

Comparators of interest include no testosterone replacement.

Outcomes

The general outcomes of interest are overall survival (OS), symptoms, morbid events, functional outcomes, and quality of life.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

A large number of systematic reviews that assess only RCTs or RCTs and non-RCTs have been published over the last 2 decades and reported on outcomes such as sexual function, (9-11) body composition, (12, 13) and bone mineral density (BMD), (14, 15) while individual RCTs have evaluated the impact of testosterone on depression (16-18) and cognition. (19) These are not

detailed here because they are outdated and/or include results of non-RCTs. A more recent systematic review that only included data from RCTs with low bias is discussed next.

The Endocrine Society commissioned a systematic review and meta-analysis, conducted by Ponce et al. (2018) to determine whether testosterone replacement therapy 1) improves sexual function, physical function, fatigue, mood, cognition, anemia, and BMD in men with hypogonadism and 2) is associated with an increased risk of lower urinary tract symptoms and erythrocytosis in men with hypogonadism. (20) The systematic review evaluated only placebo-controlled randomized trials (4 RCTs; N=1779) that assigned men with symptomatic hypogonadism with total testosterone level less than 300 ng/dL at the screening. The systematic review of characteristics and results is summarized in Tables 1 and 2. Results reported that testosterone replacement therapy was associated with a small but statistically significant improvement in libido, erectile function, sexual activity, and sexual satisfaction compared with placebo but no differences in energy levels or mood. Compared with placebo, testosterone treatment was associated with a significantly higher frequency of erythrocytosis but there was no significant difference in the change in lower urinary tract symptoms. Strengths of this review were the inclusion of only RCTs that were low-risk of bias, participants who met criteria for the diagnosis of hypogonadism (testosterone level ≤ 300 ng/dL, and presence of ≥ 1 symptoms or signs of hypogonadism), and reported outcomes were deemed clinically relevant and important to patients and ascertained using validated instruments. Limitations included the heterogeneity of instruments used to ascertain outcomes across trials, hypogonadism of multiple etiologies and lack of individual patient data meta-analysis to ascertain the relation between symptoms improvement and testosterone levels. Further, none of the trials selected in the systematic review were long enough or large enough to have sufficient statistical power to ascertain safety outcomes (prostate cancer, cardiovascular events, bone fractures).

Table 1. Systematic Review Characteristics

Study	Dates	Trials	Participants	N (Range)	Design	Duration, week
Ponce et al. (2018) (20)	To 2017	4	Symptomatic hypogonadism with total testosterone level <300 ng/dL at screening	1779 (NR)	RCT	12-52

NR: not reported; RCT: randomized controlled trial.

Table 2. Systematic Review Results

Study	Libido	Erectile Function	Sexual Activity	Sexual Satisfaction	Energy	Mood	Erythrocytosis	LUTS
Ponce et al. (2018) (20)								
n	1383	1344	1486	676	1503	1179	1579	866
SMD, RR, or MD (95% CI)	0.17 ^a (0.01 to 0.34)	0.16 ^a (0.06 to 0.27)	0.23 ^a (0.13 to 0.33)	0.16 ^a (0.01 to 0.31)	0.08 ^a (-0.02 to 0.18)	0.08 ^a (-0.03 to 0.20)	8.14 ^b (1.87 to 35.40)	0.38 ^c (20.67 to 1.43)

CI: confidence interval; LUTS: lower urinary tract symptoms; MD: mean difference; RR: relative risk; SMD: standardized mean difference.

^a SMD.

^b RR.

^c MD.

Randomized Controlled Trials

The National Institutes of Health sponsored 7 double-blind, placebo-controlled randomized trials that evaluated whether testosterone treatment of elderly men with low serum testosterone concentrations and symptoms and objective evidence of impaired mobility and/or diminished libido and/or reduced vitality would be efficacious in improving mobility (Physical Function Trial), sexual function (Sexual Function Trial), fatigue (Vitality Trial), cognitive function (Cognitive Function Trial), hemoglobin (Anemia Trial), bone density (Bone Trial), and coronary artery plaque volume (Cardiovascular Trial). (21) The major consideration in participant selection in these trials was a requirement of serum testosterone low enough to ensure that the men were unequivocally testosterone deficient, but not so low as to preclude sufficient enrollment or eventual generalizability of the results. Men were randomized to 12 months of testosterone gel (1%) or to placebo gel. General eligibility criteria for all trials included age 65 years or older with serum testosterone levels averaging less than 275 ng/dL. Exclusion criteria were a history of prostate cancer, at high-risk of prostate cancer, International Prostate Symptom Score (IPSS) greater than 19, a condition known to cause hypogonadism, or at high cardiovascular risk. There were also trial-specific eligibility criteria. The Sexual Function Trial required decreased libido and a partner willing to have intercourse twice a month. The Physical Function Trial required difficulty walking or climbing stairs and gait speed of less than 1.2 m/s on the 6-minute walk test. The Vitality Trial required low vitality (self-report and indicated by the score on a validated test). Results are summarized in Table 3. Briefly, testosterone treatment resulted in substantial benefit for sexual function, anemia, and bone density outcomes but had a small impact on physical function and vitality and no effect on cognition. Although testosterone treatment was associated with an increase in coronary artery noncalcified plaque volume, the number of cardiovascular or prostate adverse events were comparable with the placebo arm. The major limitation of these trials is that the results apply only to men ages 65 years and older with confirmed testosterone concentrations less than 275 ng/dL and durability of treatment effect beyond a year has not been demonstrated. (22)

Table 3. Results From the Primary Outcomes of the 7 NIH-Sponsored Testosterone Trials

Trial and Outcomes	N	Mean Difference or OR (95% CI)^a	Effect Size (95% CI)^b	p
Sexual Function Trial				
PDQ-Q4 score ^c	459	0.58 (0.38 to 0.78)	0.45 (0.30 to 0.60)	<0.001
Sexual desire, DISF-M-II score	470	2.93 (2.13 to 3.74)	0.44 (0.32 to 0.56)	<0.001
Erectile function, IIEF score	470	2.64 (1.68 to 3.61)	0.32 (0.20 to 0.44)	<0.001
Physical Function Trial				
Men with ≥50 m increase in 6MWD test, % ^c	387	1.42 (0.83 to 2.45)	Not reported	0.20

6MWD, m	387	4.09 (-3.00 to 11.8)	0.06 (0.004 to 0.16)	0.28
Men whose PF-10 score increased ≥ 8 , %	365	1.34 (0.90 to 2.00)	Not reported	0.15
PF-10 score	365	2.75 (0.20 to 5.29)	0.13 (0.01 to 0.26)	0.03
Vitality Trial				
Energy, increase ≥ 4 in FACIT-Fatigue score, % ^c	474	1.23 (0.83 to 1.84)	Not reported	0.30
Energy, FACIT-Fatigue score	471	1.21 (-0.04 to 2.46)	0.19 (-0.01 to 0.38)	0.03
SF-36 vitality score	404	2.41 (0.31 to 4.50)	0.18 (0.02 to 0.34)	0.03
Positive affect, PANAS score	463	0.47 (0.02 to 0.92)	0.14 (0.01 to 0.27)	0.04
Negative affect, PANAS score	463	-0.49 (-0.79 to -0.19)	-0.18 (-0.29 to -0.06)	<0.001
Depression, PHQ-9 score	464	-0.72 (-1.20 to -0.23)	-0.18 (-0.30 to -0.06)	0.004
Cognitive Function Trial				
Verbal memory; delayed paragraph recall ^c	493	-0.07 (-0.92 to 0.79)	-0.01 (-0.14 to 0.12)	0.88
Visual memory; Benton Visual Retention Test	492	-0.28 (-0.76 to 0.19)	-0.09 (-0.24 to 0.06)	0.24
Spatial ability; card rotation test	488	-0.12 (-1.89 to 1.65)	-0.01 (-0.13 to 0.11)	0.89
Executive function; Trail Making Test B-A, s ^d	490	-5.51 (-12.91 to 1.88)	-0.09 (-0.22 to 0.03)	0.14
Anemia Trial				
Hemoglobin increase from baseline ≥ 1.0 g/dL, % ^{c,e}	62	31.5 (3.7 to 277.8)	Not reported	0.002
Hemoglobin, g/dL ^e	62	0.83 (0.48 to 1.39)	1.30 (0.75 to 2.18)	0.001
Hemoglobin increase from baseline ≥ 1.0 g/dL, % ^f	64	8.2 (2.1 to 31.9)	Not reported	0.003
Hemoglobin, g/dL ^f	64	0.64 (0.12 to 1.17)	0.90 (0.17 to 1.65)	0.018
Bone Trial				
Spine trabecular BMD, % change from baseline ^c	207	6.8 (4.8 to 8.7)	0.23 (0.17 to 0.29)	0.001
Spine whole bone BMD, % change from baseline ^d	207	4.2 (3.2 to 5.3)	0.12 (0.09 to 0.15)	0.001
Hip trabecular BMD, % change from baseline	191	1.5 (0.9 to 2.0)	0.04 (0.03 to 0.06)	0.001
Hip whole bone BMD, % change from baseline	191	1.3 (0.8 to 1.7)	0.03 (0.02 to 0.04)	0.001
Cardiovascular Trial				
Noncalcified coronary artery plaque volume, ^c mm ³	138	41 (14 to 67)	0.11 (0.04 to 0.19)	0.003
Total coronary artery plaque volume, mm ³	138	47 (13 to 80)	0.09 (0.02 to 0.15)	0.006

Coronary artery calcium score, Agatston units	138	-27 (-80 to 26)	-0.03 (-0.07 to 0.02)	0.31
---	-----	-----------------	-----------------------	------

Adapted from Snyder et al. (2018). (22)

BMD: bone mineral density; CI: confidence interval; DISF-M-II: Derogatis Inventory of Sexual Function-Men-II, sexual desire domain (range, 0-33); FACIT-Fatigue: Functional Assessment of Chronic Illness Therapy-Fatigue (range, 0-52, higher scores indicate less fatigue); IIEF: International Index of Erectile Function, erectile function domain (range, 0-30); NIH: National Institutes of Health; OR: odds ratio; PANAS: Positive and Negative Affect Scale (range, 5-50); PDQ-Q4: Psychosexual Daily Questionnaire (range, 0-12, higher scores indicate a greater number of activities); PF-10, physical function scale of the Medical Outcomes Short Form Health Survey (range, 0-100); PHQ-9: Patient Health Questionnaire 9 (range, 0-27; higher scores indicate a greater degree of depressive symptoms); SF-36, 36-item Short-Form Survey (range, 0-100); 6MWD: 6-minute walk distance.

^a Treatment effect: for continuous outcomes, the treatment effect was the mean change in men allocated to testosterone minus the mean change in men allocated to placebo, adjusted for balancing factors: baseline total testosterone level (≤ 200 or > 200 ng/dL), age (≤ 75 or > 75 years), trial site, participation in the main trials, use or nonuse of antidepressants, use or nonuse of phosphodiesterase type 5 inhibitors, and baseline value of the outcome variable; for binary outcomes, the adjusted OR was the ratio of the outcome in men allocated to testosterone to the outcome in men allocated to placebo, adjusted for the same balancing factors.

^b The effect size for continuous outcomes was calculated from the mean difference divided by the baseline standard deviation pooled across treatment arms; an effect size of 0.2 is considered a small effect, 0.5 a medium effect, and 0.8 a large effect.

^c Primary outcome measure.

^d Lower scores reflect better function.

^e Among men with unexplained anemia.

^f Among men with anemia of known cause.

Section Summary: Androgen Deficiency and Clinical Symptoms of Hypogonadism

For men with low testosterone levels and sexual dysfunction, evidence from RCTs and meta-analyses has demonstrated a beneficial effect on increased libido. Other sexual symptoms (e.g., erectile dysfunction) are also likely to be improved, but the evidence is less strong. For non-sexual symptoms, there is evidence that lean body mass increased, body fat decreased, and BMD increased with testosterone therapy. However, the impact of these changes on functional status and fractures is less clear. For outcomes such as decreased energy, depression, quality of life, and cognition, the evidence is limited and inconsistent in reporting the benefits of replacement therapy.

Androgen Deficiency and Human Immunodeficiency Virus (HIV) Infection

There is a high prevalence of androgen deficiency in individuals with HIV infection who are on antiretroviral treatment, with up to 25% of this population having low testosterone levels. Men with low levels of testosterone have worse outcomes of HIV disease, including faster disease progression, greater loss of muscle mass, and larger declines in physical functioning. (2)

Clinical Context and Therapy Purpose

The purpose of testosterone replacement therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with androgen deficiency and HIV infection.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with androgen deficiency and HIV infection.

Interventions

The therapy being considered is testosterone replacement therapy as previously described.

Comparators

Comparators of interest include no testosterone replacement.

Outcomes

The general outcomes of interest are OS, symptoms, morbid events, functional outcomes, and quality of life.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

A systematic review of testosterone replacement in HIV-infected men with weight loss was performed by Bhasin et al. (2006). (23) They identified 8 trials of testosterone replacement in HIV-infected patients with weight loss. The trials were of variable quality and heterogeneous in their methodologies. Combined analysis of changes in body weight, fat-free mass, and lean body mass was performed. There was an estimated increase of 1.1 kg in body weight (95% confidence interval [CI], 0.2 to 2.0 kg), 1.4 kg in fat-free mass (95% CI, 0.7 to 2.1 kg), and 1.3 kg in lean body mass (95% CI, 0.4 to 2.2 kg) associated with testosterone replacement. Reviewers also assessed the outcomes of muscle strength and depression. Three trials reported on changes in muscle strength, with 2 of 3 reporting significant improvements with testosterone therapy. Four trials reported on changes in depression, with combined analysis showing a modest improvement in depression scores for testosterone-treated patients. There were no significant changes in parameters of HIV infection (e.g., T lymphocyte or viral load for patients treated with testosterone).

Another systematic review and meta-analysis was performed by Santi et al. (2021) to determine the effectiveness of androgen administration on body composition in HIV-infected men. (24) They identified 17 trials (N=1267) investigating testosterone replacement therapy in HIV-infected individuals with weight loss. The studies were highly heterogeneous in their methodologies. Body weight changes were reported in 12 studies (n=734). There was an estimated mean difference (MD) increase in weight of 0.99 kg (95% CI, 0.25 to 1.72; p=.008), favoring testosterone. In a similar trend, the lean body mass and fat free mass changes were higher in the testosterone group compared to the control group (lean body mass MD, 2.67; 95% CI, 1.46 to 3.87; p<.0001; fat free mass MD, 2.99; 95% CI, 2.38 to 3.60; p<.00001). These results are limited by the high heterogeneity in methods amongst included trials.

Section Summary: Androgen Deficiency and HIV Infection

RCTs of patients with HIV infection and weight loss, included in 2 systematic reviews, found that testosterone replacement was associated with an increase in body weight and lean body mass and a decrease in body fat. Findings from these trials would suggest that testosterone replacement is likely to ameliorate the weight loss associated with HIV infection.

Androgen Deficiency and Chronic Steroid Treatment

Individuals treated with chronic steroid therapy have lower levels of testosterone compared with age-matched individuals without chronic steroid use. This effect of steroids is thought to suppress the hypothalamic-pituitary axis as well as testosterone production in the testes. This hormonal suppression contributes to the increase in abdominal fat and decrease in BMD seen in individuals treated chronically with steroids.

Clinical Context and Therapy Purpose

The purpose of testosterone replacement therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with androgen deficiency on chronic steroid treatment.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with androgen deficiency on chronic steroid treatment.

Interventions

The therapy being considered is testosterone replacement therapy as previously described.

Comparators

Comparators of interest include no testosterone replacement.

Outcomes

The general outcomes of interest are OS, symptoms, morbid events, functional outcomes, and quality of life.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

The systematic review by Bhasin et al. (2006; discussed previously) identified 2 placebo-controlled randomized trials of testosterone replacement in patients on chronic steroid treatment for asthma or chronic obstructive pulmonary disease. (23) The trials were limited by small sample sizes and short duration of follow-ups. Pooled analysis of the 2 trials showed a significant increase in lean body mass of 2.3 kg (95% CI, 2.0 to 3.6 kg) and a significant decrease in fat mass of 3.1 kg (95% CI, -2.8 to -3.5 kg). There was also a significant improvement in lumbar bone density of 4% (95% CI, 2% to 7%), although there was no significant improvement in BMD of the femoral neck.

Section Summary: Androgen Deficiency and Chronic Steroid Treatment

A meta-analysis of 2 RCTs in men receiving chronic steroid treatment found a significant increase in lean body mass and a significant decrease in fat mass in individuals receiving testosterone therapy vs placebo. Thus, the evidence would suggest that testosterone is likely to ameliorate adverse events related to chronic steroid use on these parameters.

Androgen Deficiency and Type 2 Diabetes

Clinical Context and Therapy Purpose

The purpose of testosterone replacement therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with androgen deficiency and type 2 diabetes.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with androgen deficiency and type 2 diabetes.

Interventions

The therapy being considered is testosterone replacement therapy as described above.

Comparators

Comparators of interest include no testosterone replacement.

Outcomes

The general outcomes of interest are OS, symptoms, morbid events, functional outcomes, and quality of life.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

Several systematic reviews have evaluated the use of testosterone supplementation in men with type 2 diabetes and hypogonadism. Selected systematic reviews are summarized in Tables 4 to 6. Kumar et al. (2022) published a systematic review and meta-analysis evaluating testosterone supplementation with men with hypogonadism and type 2 diabetes. (25) Twelve RCTs and 1 observational study (N=1596) were included. Improved glycemic control and lipid profiles were found in men treated with testosterone.

Zhang et al. (2018) published a systematic review and meta-analysis evaluating the effects of testosterone supplement treatment in hypogonadal men with type 2 diabetes. (26) Eight RCTs with a total of 596 participants were included (all but 3 of which are also included in Cai et al. [2014] below, and all of which are included in Kumar et al. [2022] above). Meta-analysis showed that testosterone supplement treatment can significantly improve glycemic control by reducing homeostatic model assessment of insulin resistance (MD, -0.79; 95% CI, -1.23 to -0.34), fasting glucose (MD, -0.98; 95% CI, -1.13 to -0.54), fasting insulin (MD, -2.47; 95% CI, -3.99 to -0.95), and hemoglobin A_{1c} (HbA_{1c}) percentage (MD, -0.45; 95% CI, -0.73 to -0.16). Also, results showed a decline in cholesterol (MD, -0.29; 95% CI, -0.38 to -0.19) and triglycerides (MD, -0.37; 95% CI, -0.59 to -0.15). Study limitations include lack of generalizability due to the racial and ethnic homogeneity of study populations, adjusted estimates were not performed due to insufficient data, and the variation in testosterone regimens between studies.

Cai et al. (2014) reported on the results of a systematic review and meta-analysis of RCTs that evaluated the effect of testosterone therapy on metabolic parameters in patients with type 2 diabetes and hypogonadism. (27) Five RCTs (N=351) identified met eligibility criteria, 3 of which were double-blind, placebo-controlled trials and 2 of which were open-label and single-blind, no-treatment controlled trials. In pooled analysis, testosterone was associated with reduced fasting plasma glucose levels (MD, -1.10; 95% CI, -1.88 to -0.31), fasting insulin levels (MD, -2.73; 95% CI, -3.63 to -1.84), HbA_{1c} level (MD, -0.87; 95% CI, -1.32 to -0.42), and triglyceride levels (MD, -0.35; 95% CI, -0.62 to -0.07).

Table 4. Comparison of Studies Included in Systematic Reviews and Meta-Analyses

Study	Kumar et al. (2022) (25)	Zhang et al. (2018) (26)	Cai et al. (2014) (27)
Boyanov (2003) (28)	X	X	X
Dhindsa (2015) (29)	X	X	
Gianatti (2014) (30)	X	X	
Gopal (2010) (31)	X	X	X
Groti (2018) (32)	X		
Groti (2020) (33)	X		
Hackett (2014) (34)	X	X	
Hackett (2019) (35)	X		
Heufelder (2009) (36)	X	X	X
Jones (2011) (37)	X	X	X
Kapoor (2006) (38)	X	X	X
Khripun (2018) (39)	X		
Yassin (2019) (40)	X		

Table 5. Systematic Reviews and Meta-Analysis Characteristics

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Kumar et al. (2022) (25)	Through May 5, 2022	13	Men with hypogonadism and type 2 diabetes	1596 (22-537)	12 RCT; 1 Observational	NR
Zhang et al. (2018) (26)	Up to January 2018	8	Men with hypogonadism and type 2 diabetes	596 (22-186)	RCT	12-52 weeks
Cai et al. (2014) (27)	Through July 2013	5	Men with hypogonadism and type 2 diabetes	351 (22-137)	RCT	NR

NR: not reported; RCT: randomized controlled trial.

Table 6. Systematic Review and Meta-Analysis Results

Study	Fasting Glucose	HbA _{1c}	HDL	TG	BMI
Kumar et al. (2022) (25)					
Total N	1102	1454	1639	1639	1291
WMD (95% CI)	-0.35 (-0.79 to 0.10)	-0.35 (-0.64 to -.006)	0.07 (0.00 to 0.13)	-0.23 (-0.43 to -0.03)	-0.16 (-0.45 to 0.14)
I ² (p)	69.7% (.07)	47.5% (.17)	0% (.8)	79.2% (.03)	0% (.58)
Zhang et al. (2018) (26)					

WMD (95% CI)	-0.98 (-1.43 to -0.54)	-0.45 (-0.76 to -0.16)	0.01 (-0.08 to 0.09)	-0.37 (-0.59 to -0.15)	0.29 (-0.84 to 1.41)
I^2 (p)	82.7% (.00)	86.9% (.00)	88.0% (.00)	60.1% (.014)	59.9% (.021)
Cai et al. (2014) (27)					
Total N	300	124		269	
WMD (95% CI)	-1.10 (-1.88 to -0.31)	-0.87 (-1.32 to -0.42)		-0.35 (-0.62 to -0.07)	
I^2 (p)	78.7% (.009)	36% (.0001)		0% (.01)	

BMI: body mass index; CI: confidence interval; HbA_{1c}: hemoglobin A_{1c}; HDL: high-density lipoprotein; TG: triglycerides; WMD: weighted mean difference.

Section Summary: Androgen Deficiency and Type 2 Diabetes

Several systematic reviews of RCTs have assessed testosterone replacement in individuals with hypogonadism and type 2 diabetes. Beneficial effects of testosterone replacement were seen on blood glucose and lipid parameters; however, authors of the most recent systematic review noted several limitations including the short follow-up period in many studies, the variability in testosterone dose and dosage form, and lack of clinical outcomes including cardiovascular events. In addition, the majority of studies were noted to have selective reporting bias. The benefits may be outweighed by the increased risk of adverse events of treatment in the diabetic population.

Older Men with Low Testosterone Levels Without Definite Hypogonadism

Clinical Context and Therapy Purpose

The purpose of testosterone replacement therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as no testosterone replacement, in individuals who are older men with low testosterone levels without definite hypogonadism.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who are older men with low testosterone levels without definite hypogonadism.

Interventions

The therapy being considered is testosterone replacement therapy.

Comparators

Comparators of interest include no testosterone replacement.

Outcomes

The general outcomes of interest are OS, symptoms, morbid events, functional outcomes, and quality of life.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Randomized Controlled Trials

A few RCTs have evaluated the impact of testosterone replacement in elderly men with low testosterone levels, without definite evidence of hypogonadism. Most trials have been small and included only a limited range of outcomes. The largest RCTs are discussed next.

Mok et al. (2020) published the results of a randomized, double-blind, placebo-controlled study that enrolled 45 men aged at least 40 years without pathologic hypogonadism but with androgen deficiency-like energy and/or sexual symptoms to either daily testosterone or placebo gel treatment for 6 weeks. (41) The trial included 3 phases including a cross-over study design for the first 2 phases followed by a third mandatory extension phase in which participants chose which previous treatment they preferred to repeat while remaining masked to their original treatment. Primary endpoints were energy and sexual symptoms as assessed by a visual analog scale called lead symptom score. Results showed that 6 weeks of treatment with testosterone did not improve energy or sexual symptoms more than placebo in symptomatic men without pathologic hypogonadism.

Traustadottir et al. (2018) performed a double-blind, randomized, placebo-controlled, parallel-group trial to determine the effects of testosterone supplementation on oxygen consumption (VO₂) peak during incremental cycle ergometry for older men with low testosterone. (42) Patients were randomized to either the testosterone group (n=69) or placebo group (n=60). Men in the testosterone group maintained the same VO₂ peak from baseline (24.2 ± 5.2 mL/kg/min); however, the VO₂ peak fell significantly from baseline (23.6 ± 5.6 mL/kg/min) for the placebo group (average 3-year decrease, 0.88 mL/kg/min; 95% CI, -1.39 to 0.38 mL/kg/min; p=0.035). There was significant change in the difference in VO₂ peak between groups (average 3-year difference, 0.91 mL/kg/min; 95% CI, 0.010 to 0.122 mL/kg/min; p=0.008). A limitation of the study was the participants who completed measures of aerobic capacity across the time points were limited to one site.

A larger multicenter RCT reported by Legros et al. (2009) from Europe, enrolled 322 patients who were 50 years or older, with mild-to-moderate symptoms of hypogonadism and low testosterone levels. (43) Patients were randomized to daily testosterone 80, 160, or 240 mg or to placebo, and the primary outcome was the change in the Aging Males Symptom (AMS) scale at 6 months. There were no statistically significant differences in the total AMS score between groups at 6 months, although the scores in the testosterone group showed a greater numeric improvement. There was a statistically significant difference in the AMS sexual domain

subscore for the testosterone 160-mg testosterone group, but not for the 80- and the 240-mg groups. There were no statistically significant differences in adverse events between groups, including the change in prostate-specific antigen (PSA) level.

A trial by Emmelot-Vonk et al. (2008) enrolled 237 men between the ages of 60 to 80 years who had low testosterone levels but were otherwise healthy. (44) Patients were randomized to oral testosterone 80 mg or to placebo and followed for 6 months. A range of outcome measures were reported, including functional mobility, body composition, muscle strength, cognitive function, BMD, metabolic parameters, and quality of life. Safety outcomes were also included: PSA, prostate volume, renal function, liver function, and hematocrit levels. For most outcome measures, there was no improvement in the testosterone group compared with placebo therapy. There was an increase in lean body mass and a decrease in the percent body fat. However, these changes were not accompanied by improvements in functional capacity or muscle strength. There were no significant changes in cognitive function, BMD, or quality of life. There was a worsening metabolic profile, though not statistically significant ($p=0.07$), with 47.8% of men in the testosterone group meeting the definition for metabolic syndrome at the end of the study compared with 35.5% of men in the placebo group. There was a significant, but small, increase in hematocrit concentration for men in the testosterone group, and an increase in creatinine levels that was of borderline significance. Otherwise, there were no group differences in safety outcomes.

Several smaller RCTs have been published, ranging in size from 13 to 131 patients. (45-49) The most consistent finding reported in these trials was an increase in lean body mass (4 studies) and a decrease in body fat (3 studies). The impact on strength was mixed, with 2 studies reporting an improvement in the testosterone group and 2 studies reporting no difference between groups. An increase in hemoglobin and/or hematocrit concentration was reported in 1 study, and an increase in BMD was reported in another. None of these RCTs reported on functional status, quality of life, or sexual performance.

Section Summary: Older Men with Low Testosterone Levels Without Definite Hypogonadism

Several RCTs have been published, and most have been small and reported on a limited range of clinical outcomes. For most outcomes reported, there was no significant benefit reported for testosterone replacement. Some studies have reported improvements in lean body mass and decreased body fat, and 1 RCT reported improvement in sexual function. However, these trials did not report improvements in functional status or muscle strength. The adverse event profile of testosterone therapy is not well-defined, and there have been concerns about increased adverse prostate-related outcomes and cardiovascular outcomes. This uncertainty in the adverse event profile creates challenges in determining the risk-benefit profile of treatment in otherwise healthy men.

Adverse Events of Testosterone Therapy

There is a long list of potential adverse events with testosterone replacement, as follows (6):

- Prostate-related events, including development or worsening of prostate cancer, prostatic hypertrophy, increases in PSA levels, and symptoms of prostatism;

- Cardiovascular events;
- Adverse changes in lipid profile;
- Erythrocytosis and increases in hematocrit;
- Precipitation or worsening of sleep apnea;
- Liver toxicity;
- Suppression of spermatogenesis;
- Acne;
- Worsening of male pattern baldness; and
- Gynecomastia.

The clinical significance of many of these potential adverse events is unclear. Several meta-analyses have studied adverse events expected to be more common. This review of adverse events will include both randomized and nonrandomized studies, with emphasis on systematic reviews and meta-analyses of the available studies.

Cardiovascular Events

Systematic Reviews

Corona et al. (2024) conducted a systematic review and meta-analysis evaluating the cardiovascular safety of testosterone replacement therapy in men and included 106 placebo controlled RCTs. (50) The analysis found no difference between testosterone therapy and placebo when analyzing overall major adverse cardiovascular events (MACE). The single trial that looked at cardiovascular safety as the primary endpoint showed an increase in the incidence of non-fatal arrhythmias and atrial fibrillation; however, when all other trials were taken into account, this finding was not supported.

Hudson et al. (2022) evaluated the cardiovascular risk associated with testosterone use in 35 studies (N=5601) published between 1992 and August 27, 2018. (51) Risk of cardiovascular events was similar during testosterone and placebo treatment in men with hypogonadism (7.5% vs. 7.2%; odds ratio [OR], 1.07; 95% CI, 0.81 to 1.42; $p=.62$). The authors concluded that short- to medium-term studies do not indicate increased risk of cardiovascular events with testosterone replacement in men with hypogonadism.

Albert and Morley (2016) reported the findings of a systematic review that included 45 trials with 5328 subjects with mean age 63.3 years and a mean follow-up of 10.6 months. (52) Over the duration of available follow-up, testosterone treatment was not associated with increased risk of cardiovascular events (RR=1.10; 95% CI, 0.86 to 1.41; $p=0.45$). However, there was an increase event rate during the first 12 months (RR=1.79; 95% CI, 1.13 to 2.83; $p=0.012$), predominantly among those 65 years or older (RR=2.90; 95% CI, 1.35 to 6.21; $p=0.006$).

Corona et al. (2015) published an overview of previously published meta-analyses on the association between testosterone replacement therapy and cardiovascular risk. (53) Reviewers included the 3 meta-analyses described below, as well as two others, all of which focused on RCT evidence. Reviewers reported that only one of the 5 meta-analyses supported an

association between testosterone therapy and an increased cardiovascular risk. They stated that, in the single positive meta-analysis, cases of peripheral edema and self-reported syncope were included in the category of cardiovascular events, which might have overstated the number of clinically significant events. The other 4 meta-analyses did not find significant differences between testosterone and placebo groups in the incidence of overall cardiovascular events or specific events including cardiovascular death, fatal and nonfatal myocardial infarctions (MI), and cerebrovascular events.

Fernandez-Balsells et al. (2010) published a systematic review of randomized and nonrandomized comparative studies. (54) They included 51 studies and examined the outcomes of mortality, cardiovascular events, cardiovascular risk factors, prostate events, and erythrocytosis. Patients treated with testosterone had increased hematocrit concentrations (weighted mean difference [WMD], 3.2%; 95% CI, 1.4% to 5.0%), and decreased high-density lipoprotein (HDL) levels (WMD, 0.5 mg/dL; 95% CI, 0.13 to 0.85 mg/dL). No significant differences were reported in mortality, cardiovascular events, or prostate-related events.

Calof et al. (2005) performed a meta-analysis of placebo-controlled randomized trials. (55) Nineteen studies were selected (N=1084 patients) and assessed the outcomes of mortality, prostate-related events, changes in hematocrit concentration, and sleep apnea. Patients treated with testosterone were more likely to have a hematocrit greater than 50% (OR=3.7; 95% CI, 1.8 to 7.5). There were no significant differences in mortality, cardiovascular events, or sleep apnea.

In another systematic review of placebo-controlled randomized trials, Haddad et al. (2007) examined the rates of adverse cardiovascular events and changes in cardiovascular risk factors. (56) This review included 30 trials (N=1642 men). Total adverse cardiovascular events were numerically more frequent in testosterone-treated patients, but the difference compared with placebo was not statistically significant (OR=1.8; 95% CI, 0.8 to 4.2). There were small changes in blood pressure, lipid levels, and glucose, but none of these changes were statistically significant.

Randomized Controlled Trials

Lincoff et al. (2023) evaluated the cardiovascular safety of testosterone replacement therapy in a multicenter noninferiority trial in middle-aged and older men with hypogonadism (TREAVERSE trial). (57) Patients (N=5,246) were randomized to receive daily transdermal testosterone gel or placebo gel. The noninferiority margin was an upper limit of <1.5 for the 95% CI of the hazard ratio among patients who received at least 1 dose of testosterone or placebo. One hundred eighty-two patients (7.0%) in the testosterone group and 190 patients (7.3%) in the placebo group experienced a primary cardiovascular outcome incident (hazard ratio, 0.96; 95% CI, 0.78 to 1.17; $p < 0.001$ for noninferiority). In both groups, the incidence of secondary end-point events (i.e., first occurrence of any component of the composite of death from cardiovascular causes, nonfatal myocardial infarction, nonfatal stroke, or coronary revascularization in a time-to-event analysis) were comparable. This was the trial that the updated 2025 FDA Drug Safety Communication labeling changes were based on.

Cohort Studies

In addition to RCTs and meta-analyses of them, several large cohort studies have been published. Finkle et al. (2014) conducted a large retrospective cohort study using administrative claims data to assess the relation between testosterone therapy and nonfatal MI. (58) The authors generated a cohort of 55,593 men who filled their first prescription for one of several testosterone prescriptions between 2008 and 2010 from the Truven Health MarketScan® Commercial Claims and Encounters Database, which includes diagnoses, procedures, and prescriptions for all enrollees of contributing health plans. Testosterone recipients were compared with a population of men who filled their first prescription for a phosphodiesterase type 5 inhibitor (PDE5I; sildenafil or tadalafil; n=167,279) during the same time period. For testosterone recipients, the rate ratio (RR) of MI in the post- compared with the pre-testosterone period was 1.36 (95% CI, 1.03 to 1.81). Compared with subjects in the PDE5I group, the rate ratio for MI risk for testosterone recipients was 1.90 (95% CI, 1.04 to 3.49). After stratifying by age, for testosterone recipients younger than age 55 years, the RR for MI in the post-testosterone period was 0.95 (95% CI, 0.54 to 1.67); for testosterone recipients ages 75 and older, the RR for MI in the post-testosterone period was 3.43 (95% CI, 1.54 to 7.56; p=0.03 for trend). No similar trend was seen for PDE5I recipients. Although this study suggested an association between testosterone use and nonfatal MI, it was limited by its retrospective design and the potential for confounding by measured and unmeasured variables.

Another large retrospective cohort study conducted by Baillargeon et al. (2014) evaluated administrative claims data from Medicare to assess the relation between testosterone administered IM and the risk of MI. (59) The study included 6355 Medicare beneficiaries who received at least 1 testosterone injection between 1997 and 2005 and who were matched in a 1:3 ratio to 19,065 testosterone nonusers based on a composite MI prognostic score. After adjustment for demographic and clinical covariates, testosterone therapy was not associated with an increased risk of MI (adjusted hazard ratio [HR]=0.84; 95% CI, 0.69 to 1.01). Testosterone therapy was associated with a reduced risk of MI in men with a MI prognostic score in the highest quartile (HR=0.69; 95% CI, 0.53 to 0.92), while men in the lower three quartiles showed no difference in MI risk with testosterone therapy.

A large retrospective comparative cohort study by Vigen et al. (2013) evaluated the risk of cardiovascular events in patients treated with testosterone replacement therapy. (60) This study used data from the Veterans Administration Clinical Assessment Reporting and Tracking Program to identify all male patients who had both undergone coronary angiography and had a total testosterone level checked between 2005 and 2011. There were 8709 patients with a low testosterone level, defined as less than 300 ng/dL. The population had high levels of comorbidity, with 80% of patients having coronary artery disease, 50% diabetes, and 20% prior to MI. There were 1223 patients treated with testosterone and 7486 who were not. After a mean follow-up of 27.5 months, the primary outcome of all-cause mortality, MI, or stroke was more frequent in the group treated with testosterone (HR, 1.29; 95% CI, 1.04 to 1.58; p=0.02).

A case-control study performed within a cohort of 934,283 men ages 45 to 80 years was published by Etminan et al. (2015). (61) It identified 30,066 cases of MI and matched each case with 4 controls. There was no evidence of increased current testosterone replacement therapy use in case patients (RR, 1.01; 95% CI, 0.89 to 1.16). There was also no association between past testosterone replacement therapy use and MI or evidence of different risk level by type of preparation. A small increase in risk was reported for first-time testosterone replacement therapy users (RR, 1.41; 95% CI, 1.06 to 1.87).

Effects on Prostate Gland

Several meta-analyses have specifically evaluated the relation between testosterone and prostate-related events. Cui et al. (2013) conducted systematic review of RCTs, which reported on the effect of testosterone replacement on prostate growth. (62) Sixteen RCTs comparing testosterone replacement with placebo (n=1030 patients) were included, 7 of which were short term (<12 months) and 9 long-term (12 to 36 months). Seven studies evaluated transdermally administered testosterone, while 6 evaluated injected testosterone and 3 evaluated orally administered testosterone. In the short-term, transdermal, but not orally administered or injected, testosterone administration was significantly associated with changes in PSA levels (standardized mean difference [SMD], 0.30; 95% CI, 0.07 to 0.54; p=0.002). However, there was no significant association between testosterone administration and PSA levels over the longer term. Testosterone administration by any method was not associated with significant differences in International Prostate Symptom Score (IPSS), prostate volume, or maximum urine flow rate.

In a separate publication, Cui et al. (2014) conducted a systematic review of RCTs that reported on the effect of testosterone replacement therapy on prostate cancer risk. (63) This analysis included 22 RCTs (N=2351 patients), 11 of which reported short-term (<12 months) outcomes and 11 of which reported long-term (12 to 36 months) outcomes. Five studies evaluated injectable testosterone, 1 evaluated oral testosterone, and 5 studies evaluated transdermal testosterone over the short term; there was no significant association between any administration method and prostate cancer, prostate biopsy, or prostate nodules. However, for the studies evaluating transdermal testosterone, there was a significant association between testosterone treatment and change in PSA level (SMD=0.33; 95% CI, 0.21 to 0.45; p<0.000). There was no association between testosterone therapy and abnormal PSA levels. For long-term administration, 3 studies evaluated injectable testosterone, 2 studies evaluated oral testosterone, and 6 studies evaluated transdermal administered testosterone. There was no significant association between testosterone administration by any method and prostate cancer, prostate biopsy, or prostate nodules. No significant association was found between testosterone administration over the long-term and change in PSA level.

A systematic review by Kohn et al. (2016) identified 14 RCTs on testosterone therapy in aging men and assessed lower urinary tract symptoms with the IPSS. (64) In a pooled analysis, there was no statistically significant difference in change in IPSS among men treated with testosterone vs placebo (p=0.11). Similarly, another systematic review, Kathrins et al. (2016) identified 35 prospective studies and found that most trials did not demonstrate a correlation

between testosterone therapy and enlarged prostate volume, de novo lower urinary tract symptoms, or worsening lower urinary tract symptoms. (65)

Venous Thromboembolism

A meta-analysis by Cannarella et al. (2023) evaluated the risk of testosterone replacement therapy and thromboembolic events in patients with pre-treatment total testosterone <12 nmol L⁻¹. (66) Twenty-four studies (which included 14 RCTs) were analyzed. Based on data gathered from RCTs, testosterone replacement therapy did not influence the risk of stroke (OR, 1.34; 95% CI, 0.09 to 18.97; p=0.83), arterial thrombosis (OR, 1.27; 95% CI, 0.47 to 3.43; p=0.64), MI (OR, 0.51; 95% CI, 0.11 to 2.31; p=0.39), pulmonary embolism (OR, 1.38; 95% CI, 0.27 to 7.04; p=0.70), or mortality (OR, 0.70; 95% CI, 0.20 to 2.38; p=0.56). When only evaluating observational cohort study data, there was a significant decrease in the risk of MI, venous thromboembolism, mortality, and arterial thrombotic events.

Houghton et al. (2018) published a systematic review and meta-analysis to determine if there is an association between exogenous testosterone (any route) and venous thromboembolism. (67) Eleven studies (6 RCTs, 5 observational studies) with a total of 1,251,876 (RCT, n=2236; observational, n=1,249,640) patients were included. Meta-analysis of all studies found no significant association between venous thromboembolism and testosterone therapy (OR, 1.41, 95% CI, 0.96 to 2.07); there was also no significant association when stratified by study design: RCT (OR, 2.05, 95% CI, 0.78 to 5.39), cohort studies (OR, 1.06, 95% CI, 0.85 to 1.33), and case-control studies (OR, 1.34, 95% CI, 0.78 to 2.28). The analysis was limited by high heterogeneity among the included studies ($I^2=84.4\%$).

A large case-control study by Baillargeon et al. (2015) evaluated the risk of venous thromboembolism associated with testosterone replacement therapy. (68) This study assessed 30,572 men ages 40 years and older. Subjects had a diagnosis of venous thromboembolism and were on an anticoagulant drug. Cases were matched with 3 controls on age, time of onset, location, diagnosis of hypogonadism, and the presence of a prothrombotic condition. There was no increased risk of testosterone replacement therapy in the venous thromboembolism group (OR=0.91; 95% CI, 0.38 to 2.16). The lack of an association persisted when different time frames of testosterone replacement therapy exposure were examined.

Summary of Evidence

For individuals who have androgen deficiency and clinical symptoms of hypogonadism who receive testosterone replacement therapy, the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are overall survival (OS), symptoms, morbid events, functional outcomes, and quality of life. For men with low testosterone levels and sexual dysfunction, the evidence has been fairly consistent in demonstrating a beneficial effect on increased libido. Other sexual function symptoms (e.g., erectile dysfunction) are also likely to be improved, but the evidence is less strong. For other symptoms, there is evidence that lean body mass is increased, body fat is decreased, and bone mineral density is increased with testosterone therapy. However, the impact of these changes on functional status and fractures is less clear. For outcomes such as decreased energy, depression, quality of life, and cognition,

the evidence is limited and inconsistent in reporting benefits of replacement therapy. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have androgen deficiency and human immunodeficiency virus (HIV) infection who receive testosterone replacement therapy, the evidence includes RCTs and systematic reviews. Relevant outcomes are OS, symptoms, morbid events, functional outcomes, and quality of life. A limited number of trials have included patients with HIV infection and weight loss. These trials have reported improvements in body weight, lean body mass, and a decrease in body fat, which indicates that testosterone replacement is likely to ameliorate weight loss associated with HIV infection. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have androgen deficiency on chronic steroid treatment who receive testosterone replacement therapy, the evidence includes RCTs and systematic reviews. Relevant outcomes are OS, symptoms, morbid events, functional outcomes, and quality of life. A limited number of trials have included patients with androgen deficiency in chronic steroid treatment. These trials have reported improvements in body weight, lean body mass, and a decrease in body fat, which are likely to ameliorate the effects of chronic steroids on these parameters. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have androgen deficiency and type 2 diabetes who receive testosterone replacement therapy, the evidence includes systematic reviews of RCTs. Relevant outcomes are OS, symptoms, morbid events, functional outcomes, and quality of life. The available systematic reviews have reported that testosterone replacement leads to modest improvements in glucose control (e.g., hemoglobin A_{1c} levels, insulin sensitivity) and lipid parameters. There is a lack of trials reporting on clinical outcomes, and the small benefits may be outweighed by the adverse events of treatment. Current professional guidelines reflect the controversy regarding the balance of risks and benefits. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals (older men) with low testosterone levels without definite hypogonadism who receive testosterone replacement therapy, the evidence includes RCTs and systematic reviews. Relevant outcomes are OS, symptoms, morbid events, functional outcomes, and quality of life. The available RCTs are mostly small and have reported on a limited range of clinical outcomes. For most outcomes, there was no benefit for testosterone replacement. Some studies have reported improvements in lean body mass and decreased body fat, and a recent RCT found improved sexual function. However, these studies did not report improvements in functional status or muscle strength. Although the adverse event profile of testosterone therapy is not well-defined, there are concerns about increased adverse prostate-related outcomes and cardiovascular outcomes. This uncertainty in the adverse event profile creates challenges in determining the risk-benefit profile of treatment in otherwise healthy men. The evidence is

insufficient to determine that the technology results in an improvement in the net health outcome.

Practice Guidelines and Position Statements

American Diabetes Association (ADA)

The ADA Comprehensive Medical Evaluation and Assessment of Comorbidities: *Standards of Medical Care in Diabetes-2022* recommends screening with a morning serum testosterone level for men with diabetes who have symptoms of low testosterone (hypogonadism), such as decreased sexual desire or activity, or erectile dysfunction. (69) Treatment in asymptomatic men is controversial. Testosterone replacement in men with symptomatic hypogonadism may have benefits including improved sexual function, well-being, muscle mass and strength, and bone density. In men with diabetes who have symptoms or signs of low testosterone (hypogonadism), a morning total testosterone level should be measured using an accurate and reliable assay. In men who have total testosterone levels close to the lower limit, it is reasonable to check sex hormone–binding globulin, as it is often low in diabetes and associated with lower testosterone levels. Further testing (such as luteinizing hormone and follicle-stimulating hormone levels) may be needed to determine if the patient has hypogonadism. Testosterone replacement in older men with hypogonadism has been associated with increased coronary artery plaque volume and, in some studies, an increase in cardiovascular events, which should be considered when assessing the risks and benefits of treatment. The 2023, 2024, and 2025 ADA standards of care echo the 2022 recommendations of screening with a morning serum testosterone level in men with diabetes who have symptoms or signs of hypogonadism. (70-72)

American Urological Association

The 2018 guidelines for the evaluation and management of testosterone deficiency from the American Urological Association are summarized in Table 7. The guidelines were reviewed, and validity was confirmed in 2024. (73)

Table 7. Practice Guidelines for Evaluation and Management of Testosterone Deficiency

Recommendation	SOR	LOE
Clinicians should adjust testosterone therapy dosing to achieve a total testosterone level in the middle tertial of the normal reference range.	Conditional	Grade C
Exogenous testosterone therapy should not be prescribed to men who are currently trying to conceive.	Strong	Grade A
Clinicians should not prescribe alkylated oral testosterone.	Moderate	Grade B
Commercially manufactured testosterone products should be prescribed rather than compounded testosterone, when possible.	Conditional	Grade C

SOR: strength of recommendation; LOE: level of evidence.

Endocrine Society

In 2018, the Endocrine Society published clinical practice guidelines on testosterone therapy in men with androgen deficiency (see Table 8). (1)

Table 8. Practice Guidelines on Testosterone Therapy in Men with Hypogonadism

Recommendations	SOR	QOE
"We recommend testosterone therapy in hypogonadal men to induce and maintain secondary sex characteristics and correct symptoms of testosterone deficiency."	Recommend	Moderate
"We recommend against testosterone therapy in men planning fertility in the near term or in men with breast or prostate cancer, a palpable prostate nodule or induration, a prostate-specific antigen level > 4 ng/mL, a prostate-specific antigen level > 3 ng/mL combined with a high risk of prostate cancer (without further urological evaluation), elevated hematocrit, untreated severe obstructive sleep apnea, severe lower urinary tract symptoms, uncontrolled heart failure, myocardial infarction or stroke within the last 6 months, or thrombophilia."	Recommend	Low
"In hypogonadal men 55 to 69 years old, who are being considered for testosterone therapy and have a life expectancy >10 years, we suggest discussing the potential benefits and risks of evaluating prostate cancer risk and prostate monitoring and engaging the patient in shared decision-making regarding prostate cancer monitoring. For patients who choose monitoring, clinicians should assess prostate cancer risk before starting testosterone treatment and 3 to 12 months after starting testosterone."	Suggest	Very Low
"In hypogonadal men being considered for testosterone therapy who are 40 to 69 years old and at increased risk of prostate cancer (e.g., African Americans and men with a first-degree relative with diagnosed prostate cancer), we suggest discussing prostate cancer risk with the patient and offering monitoring options."	Suggest	Very Low
"In men with type 2 diabetes mellitus who have low testosterone concentrations, we recommend against testosterone therapy as a means of improving glycemic control."	Recommend	Low

Adapted from Bhasin et al. (2018). (1)

SOR: strength of recommendation; QOE: quality of evidence.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this policy are listed in Table 9.

Table 9. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date

Ongoing			
NCT04731376	Perioperative Testosterone Replacement Therapy Improves Outcomes: A Pilot Safety and Feasibility Study	100	Dec 2026
NCT04049331	Improving Patient-Important Outcomes With Testosterone Replacement in Hypogonadal Men With a Prior History of Cancer	240	Jan 2027
NCT04301765	Improving Cancer-related Fatigue, Sexual Dysfunction and Quality of Life in Older Men With Cancer and Androgen Deficiency	230	Jan 2026
Unpublished			
NCT04456296 ^a	n Open-Label, Randomized, Parallel-Group, Three Treatment Arm, Multicenter Study on Hypogonadal Males to Evaluate the Effect on 24-Hour Ambulatory Blood Pressures After 16-Week Continuous Administration With Marketed Testosterone Products	676	Jul 2023
NCT03339635	Short-term Testosterone Replacement in Testicular Cancer Survivors to Treat Overweight and Improve Cardiometabolic Risk: A Pilot Study	32	Jun 2024

NCT: national clinical trial.

^a Denotes sponsorship or cosponsorship by the manufacturer.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member's benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

CPT Codes	11980, 11981, 11982, 11983, 96372
HCPCS Codes	G0516, G0517, G0518, J1071, J1072, J1073, J2320, J3121, J3145, [Deleted 1/2026: S0189]

*Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.

References

1. Bhasin S, Brito JP, Cunningham GR, et al. Testosterone therapy in men with hypogonadism: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* May 01 2018; 103(5):1715-1744. PMID 29562364
2. Bhasin S, Cunningham GR, Hayes FJ, et al. Testosterone therapy in men with androgen deficiency syndromes: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* Jun 2010; 95(6):2536-2559. PMID 20525905
3. Zarotsky V, Huang MY, Carman W, et al. Systematic literature review of the risk factors, comorbidities, and consequences of hypogonadism in men. *Andrology.* Nov 2014; 2(6):819-834. PMID 25269643
4. Wu FC, Tajar A, Pye SR, et al. Hypothalamic-pituitary-testicular axis disruptions in older men are differentially linked to age and modifiable risk factors: the European Male Aging Study. *J Clin Endocrinol Metab.* Jul 2008; 93(7):2737-2745. PMID 18270261
5. Travison TG, Araujo AB, Kupelian V, et al. The relative contributions of aging, health, and lifestyle factors to serum testosterone decline in men. *J Clin Endocrinol Metab.* Feb 2007; 92(2):549-555. PMID 17148559
6. Cunningham GR, Toma SM. Clinical review: Why is androgen replacement in males controversial? *J Clin Endocrinol Metab.* Jan 2011; 96(1):38-52. PMID 20881265
7. FDA Drug Safety Communication: FDA cautions about using testosterone products for low testosterone due to aging; requires labeling change to inform of possible increased risk of heart attack and stroke with use. Updated February 25, 2025. Available at <<https://www.fda.gov>> (accessed May 16, 2025).
8. FDA Drug Safety Communication: FDA issues class-wide labeling changes for testosterone products. Updated February 28, 2025. Available at <<https://www.fda.gov>> (accessed May 15, 2025).
9. Boloña ER, Uraga MV, Haddad RM, et al. Testosterone use in men with sexual dysfunction: a systematic review and meta-analysis of randomized placebo-controlled trials. *Mayo Clinic Proc.* Jan 2007; 82(1):20-28. PMID 17285782
10. Jain P, Rademaker AW, McVary KT. Testosterone supplementation for erectile dysfunction: results of a meta-analysis. *J Urol.* Aug 2000; 164(2):371-375. PMID 10893588
11. Corona G, Rastrelli G, Morgentaler A, et al. Meta-analysis of results of testosterone therapy on sexual function based on International Index of Erectile Function Scores. *Eur Urol.* Dec 2017; 72(6):1000-1011. PMID 28434676
12. Neto WK, Gama EF, Rocha LY, et al. Effects of testosterone on lean mass gain in elderly men: systematic review with meta-analysis of controlled and randomized studies. *Age (Dordr).* Feb 2015; 37(1):9742. PMID 25637335
13. Bhasin S, Calof OM, Storer TW, et al. Drug insight: Testosterone and selective androgen receptor modulators as anabolic therapies for chronic illness and aging. *Nat Clin Pract Endocrinol Metab.* Mar 2006; 2(3):146-159. PMID 16932274
14. Corona G, Vena W, Pizzocaro A, et al. Testosterone supplementation and bone parameters: a systematic review and meta-analysis study. *J Endocrinol Invest.* May 2022; 45(5):911-926. PMID 35041193
15. Tracz MJ, Sideras K, Boloña ER, et al. Testosterone use in men and its effects on bone health. A systematic review and meta-analysis of randomized placebo-controlled trials. *J Clin Endocrinol Metab.* Jun 2006; 91(6):2011-2016. PMID 16720668

16. Amiaz R, Pope HG, Mahne T, et al. Testosterone gel replacement improves sexual function in depressed men taking serotonergic antidepressants: a randomized, placebo-controlled clinical trial. *J Sex Marital Ther.* 2011; 37(4):243-254. PMID 21707327
17. Shores MM, Kivlahan DR, Sadak TI, et al. A randomized, double-blind, placebo-controlled study of testosterone treatment in hypogonadal older men with subthreshold depression (dysthymia or minor depression). *J Clin Psychiatry.* Jul 2009; 70(7):1009-1016. PMID 19653976
18. Seidman SN, Orr G, Raviv G, et al. Effects of testosterone replacement in middle-aged men with dysthymia: a randomized, placebo-controlled clinical trial. *J Clin Psychopharmacol.* Jun 2009; 29(3):216-221. PMID 19440073
19. Borst SE, Yarrow JF, Fernandez C, et al. Cognitive effects of testosterone and finasteride administration in older hypogonadal men. *Clin Interv Aging.* 2014; 9:1327-1333. PMID 25143719
20. Ponce OJ, Spencer-Bonilla G, Alvarez-Villalobos N, et al. The efficacy and adverse events of testosterone replacement therapy in hypogonadal men: A systematic review and meta-analysis of randomized, placebo-controlled trials. *J Clin Endocrinol Metab.* Mar 17 2018. PMID 29562341
21. Snyder PJ, Ellenberg SS, Cunningham GR, et al. The testosterone trials: seven coordinated trials of testosterone treatment in elderly men. *Clin Trials.* Jun 2014; 11(3):362-375. PMID 24686158
22. Snyder PJ, Bhasin S, Cunningham GR, et al. Lessons from the testosterone trials. *Endocr Rev.* Jun 01 2018; 39(3):369-386. PMID 29522088
23. Bhasin S, Cunningham GR, Hayes FJ, et al. Testosterone therapy in adult men with androgen deficiency syndromes: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab.* Jun 2006; 91(6):1995-2010. PMID 16720669
24. Santi D, Spaggiari G, Vena W, et al. The Prevalence of Hypogonadism and the Effectiveness of Androgen Administration on Body Composition in HIV-Infected Men: A Meta-Analysis. *Cells.* Aug 12 2021; 10(8):2067. PMID 34440836
25. Kumar S, Khatri M, Memon RA, et al. Effects of testosterone therapy in adult males with hypogonadism and T2DM: A meta-analysis and systematic review. *Diabetes Metab Syndr.* Aug 2022; 16(8):102588. PMID 35952509
26. Zhang J, Yang B, Xiao W, et al. Effects of testosterone supplement treatment in hypogonadal adult males with T2DM: a meta-analysis and systematic review. *World J Urol.* Aug 2018; 36(8):1315-1326. PMID 29511802
27. Cai X, Tian Y, Wu T, et al. Metabolic effects of testosterone replacement therapy on hypogonadal men with type 2 diabetes mellitus: a systematic review and meta-analysis of randomized controlled trials. *Asian J Andro.* 2014; 16(1):146-152. PMID 24369149
28. Boyanov MA, Boneva Z, Christov VG. Testosterone supplementation in men with type 2 diabetes, visceral obesity and partial androgen deficiency. *Aging Male.* Mar 2003; 6(1):1-7. PMID 12809074
29. Dhindsa S, Ghanim H, Batra M, et al. Insulin Resistance and Inflammation in Hypogonadotropic Hypogonadism and Their Reduction After Testosterone Replacement in Men With Type 2 Diabetes. *Diabetes Care.* Jan 2016; 39(1):82-91. PMID 26622051

30. Gianatti EJ, Dupuis P, Hoermann R, et al. Effect of testosterone treatment on glucose metabolism in men with type 2 diabetes: a randomized controlled trial. *Diabetes Care*. Aug 2014; 37(8):2098-2107. PMID 24804695
31. Gopal RA, Bothra N, Acharya SV, et al. Treatment of hypogonadism with testosterone in patients with type 2 diabetes mellitus. *Endocr Pract*. 2010; 16(4):570-576. PMID 20150021
32. Groti K, Žuran I, Antoniĉ B, et al. The impact of testosterone replacement therapy on glycemic control, vascular function, and components of the metabolic syndrome in obese hypogonadal men with type 2 diabetes. *Aging Male*. Sep 2018; 21(3):158-169. PMID 29708829
33. Groti Antoniĉ K, Antoniĉ B, Žuran I, et al. Testosterone treatment longer than 1 year shows more effects on functional hypogonadism and related metabolic, vascular, diabetic and obesity parameters (results of the 2-year clinical trial). *Aging Male*. Dec 2020; 23(5):1442-1454. PMID 32844712
34. Hackett G, Cole N, Bhartia M, et al. Testosterone replacement therapy improves metabolic parameters in hypogonadal men with type 2 diabetes but not in men with coexisting depression: the BLAST study. *J Sex Med*. Mar 2014; 11(3):840-856. PMID 24308723
35. Hackett G, Cole N, Mulay A, et al. Long-term testosterone therapy in type 2 diabetes is associated with reduced mortality without improvement in conventional cardiovascular risk factors. *BJU Int*. Mar 2019; 123(3):519-529. PMID 30216622
36. Heufelder AE, Saad F, Bunck MC, et al. Fifty-two-week treatment with diet and exercise plus transdermal testosterone reverses the metabolic syndrome and improves glycemic control in men with newly diagnosed type 2 diabetes and subnormal plasma testosterone. *J Androl*. 2009; 30(6):726-733. PMID 19578132
37. Jones TH, Arver S, Behre HM, et al. Testosterone replacement in hypogonadal men with type 2 diabetes and/or metabolic syndrome (the TIMES2 study). *Diabetes Care*. Apr 2011; 34(4):828-837. PMID 21386088
38. Kapoor D, Goodwin E, Channer KS, et al. Testosterone replacement therapy improves insulin resistance, glycemic control, visceral adiposity and hypercholesterolemia in hypogonadal men with type 2 diabetes. *Eur J Endocrinol*. Jun 2006; 154(6):899-906. PMID 16728551
39. Khripun I, Vorobyev S, Belousov I, et al. Influence of testosterone substitution on glycemic control and endothelial markers in men with newly diagnosed functional hypogonadism and type 2 diabetes mellitus: a randomized controlled trial. *Aging Male*. Dec 2019; 22(4):241-249. PMID 30235049
40. Yassin A, Haider A, Haider KS, et al. Testosterone Therapy in Men With Hypogonadism Prevents Progression From Prediabetes to Type 2 Diabetes: Eight-Year Data From a Registry Study. *Diabetes Care*. Jun 2019; 42(6):1104-1111. PMID 30862651
41. Mok SF, Fennell C, Savkovic S, et al. Testosterone for Androgen Deficiency-Like Symptoms in Men Without Pathologic Hypogonadism: A Randomized, Placebo-Controlled Cross-over With Masked Choice Extension Clinical Trial. *J Gerontol A Biol Sci Med Sci*. Sep 16 2020; 75(9):1723-1731. PMID 31425577
42. Traustadóttir T, Harman SM, Tsitouras P, et al. Long-Term Testosterone Supplementation in Older Men Attenuates Age-Related Decline in Aerobic Capacity. *J Clin Endocrinol Metab*. Aug 01 2018; 103(8):2861-2869. PMID 29846604

43. Legros JJ, Meuleman EJ, Elbers JM, et al. Oral testosterone replacement in symptomatic late-onset hypogonadism: effects on rating scales and general safety in a randomized, placebo-controlled study. *Eur J Endocrinol*. May 2009; 160(5):821-831. PMID 19211706
44. Emmelot-Vonk MH, Verhaar HJ, Nakhai Pour HR, et al. Effect of testosterone supplementation on functional mobility, cognition, and other parameters in older men: a randomized controlled trial. *JAMA*. Jan 02 2008; 299(1):39-52. PMID 18167405
45. Kenny AM, Kleppinger A, Annis K, et al. Effects of transdermal testosterone on bone and muscle in older men with low bioavailable testosterone levels, low bone mass, and physical frailty. *J Am Geriatr Soc*. Jun 2010; 58(6):1134-1143. PMID 20722847
46. Kenny AM, Prestwood KM, Gruman CA, et al. Effects of transdermal testosterone on bone and muscle in older men with low bioavailable testosterone levels. *J Gerontol A Biol Sci Med Sci*. May 2001; 56(5):M266-272. PMID 11320105
47. Snyder PJ, Peachey H, Hannoush P, et al. Effect of testosterone treatment on body composition and muscle strength in men over 65 years of age. *J Clin Endocrinol Metab*. Aug 1999; 84(8):2647-2653. PMID 10443654
48. Sih R, Morley JE, Kaiser FE, et al. Testosterone replacement in older hypogonadal men: a 12-month randomized controlled trial. *J Clin Endocrinol Metab*. Jun 1997; 82(6):1661-1667. PMID 9177359
49. Tenover JS. Effects of testosterone supplementation in the aging male. *J Clin Endocrinol Metab*. Oct 1992; 75(4):1092-1098. PMID 1400877
50. Corona G, Rastrelli G, Sparano C, et al. Cardiovascular safety of testosterone replacement therapy in men: an updated systematic review and meta-analysis. *Expert Opin Drug Saf*. May 2024; 23(5):565-579. PMID 38553429
51. Hudson J, Cruickshank M, Quinton R, et al. Adverse cardiovascular events and mortality in men during testosterone treatment: an individual patient and aggregate data meta-analysis. *Lancet Healthy Longev*. Jun 2022; 3(6):e381-e393. PMID 35711614
52. Albert SG, Morley JE. Testosterone therapy, association with age, initiation and mode of therapy with cardiovascular events: a systematic review. *Clin Endocrinol (Oxf)*. Sep 2016; 85(3):436-443. PMID 27124404
53. Corona GG, Rastrelli G, Maseroli E, et al. Testosterone replacement therapy and cardiovascular risk: a review. *World J Mens Health*. Dec 2015; 33(3):130-142. PMID 26770933
54. Fernandez-Balsells MM, Murad MH, Lane M, et al. Clinical review 1: Adverse effects of testosterone therapy in adult men: a systematic review and meta-analysis. *J Clin Endocrinol Metab*. Jun 2010; 95(6):2560-2575. PMID 20525906
55. Calof OM, Singh AB, Lee ML, et al. Adverse events associated with testosterone replacement in middle-aged and older men: a meta-analysis of randomized, placebo-controlled trials. *J Gerontol A Biol Sci Med Sci*. Nov 2005; 60(11):1451-1457. PMID 16339333
56. Haddad RM, Kennedy CC, Caples SM, et al. Testosterone and cardiovascular risk in men: a systematic review and meta-analysis of randomized placebo-controlled trials. *Mayo Clinic Proc*. Jan 2007; 82(1):29-39. PMID 17285783
57. Lincoff AM, Bhasin S, Flevaris P, et al. Cardiovascular Safety of Testosterone-Replacement Therapy. *N Engl J Med*. Jul 13 2023; 389(2):107-117. PMID 37326322

58. Finkle WD, Greenland S, Ridgeway GK, et al. Increased risk of non-fatal myocardial infarction following testosterone therapy prescription in men. *PLoS One*. 2014; 9(1):e85805. PMID 24489673
59. Baillargeon J, Urban RJ, Kuo YF, et al. Risk of Myocardial Infarction in Older Men Receiving Testosterone Therapy. *Ann Pharmacother*. Sep 2014; 48(9):1138-1144. PMID 24989174
60. Vigen R, O'Donnell CI, Baron AE, et al. Association of testosterone therapy with mortality, myocardial infarction, and stroke in men with low testosterone levels. *JAMA*. Nov 6 2013; 310(17):1829-1836. PMID 24193080
61. Etminan M, Skeldon SC, Goldenberg SL, et al. Testosterone therapy and risk of myocardial infarction: a pharmacoepidemiologic study. *Pharmacotherapy*. Jan 2015; 35(1):72-78. PMID 25582846
62. Cui Y, Zhang Y. The effect of androgen-replacement therapy on prostate growth: a systematic review and meta-analysis. *Eur Urol*. Nov 2013; 64(5):811-822. PMID 23567065
63. Cui Y, Zong H, Yan H, et al. The effect of testosterone replacement therapy on prostate cancer: a systematic review and meta-analysis. *Prostate Cancer Prostatic Dis*. Jun 2014; 17(2):132-143. PMID 24445948
64. Kohn TP, Mata DA, Ramasamy R, et al. Effects of testosterone replacement therapy on lower urinary tract symptoms: a systematic review and meta-analysis. *Eur Urol*. Jun 2016; 69(6):1083-1090. PMID 26874809
65. Kathrins M, Doersch K, Nimeh T, et al. The relationship between testosterone-replacement therapy and lower urinary tract symptoms: a systematic review. *Urology*. Feb 2016; 88:22-32. PMID 26616095
66. Cannarella R, Gusmano C, Leanza C, et al. Testosterone replacement therapy and vascular thromboembolic events: a systematic review and meta-analysis. *Asian J Androl*. Mar 01 2024; 26(2):144-154. PMID 37921515
67. Houghton DE, Alsawas M, Barrioneuvo P, et al. Testosterone therapy and venous thromboembolism: A systematic review and meta-analysis. *Thromb Res*. Dec 2018; 172:94-103. PMID 30396049
68. Baillargeon J, Urban RJ, Morgentaler A, et al. Risk of venous thromboembolism in men receiving testosterone therapy. *Mayo Clin Proc*. Aug 2015; 90(8):1038-1045. PMID 26205547
69. Draznin B, Aroda VR, Bakris G, et al. 4. Comprehensive Medical Evaluation and Assessment of Comorbidities: Standards of Medical Care in Diabetes-2022. *Diabetes Care*. Jan 01 2022; 45(Suppl 1):S46-S59. PMID 34964869
70. ElSayed NA, McCoy RG, Aleppo G, et al. 4. Comprehensive Medical Evaluation and Assessment of Comorbidities: Standards of Care in Diabetes-2025. *Diabetes Care*. Jan 01 2025; 48(1 Suppl 1):S59-S85. PMID 39651988
71. ElSayed NA, Aleppo G, Bannuru RR, et al. 4. Comprehensive Medical Evaluation and Assessment of Comorbidities: Standards of Care in Diabetes-2024. *Diabetes Care*. Jan 01 2024; 47(Suppl 1):S52-S76. PMID 38078591
72. ElSayed NA, Aleppo G, Aroda VR, et al. 4. Comprehensive Medical Evaluation and Assessment of Comorbidities: Standards of Care in Diabetes-2023. *Diabetes Care*. Jan 01 2023; 46(Suppl 1):S49-S67. PMID 36507651

73. Mulhall JP, Trost LW, Brannigan RE, et al. Evaluation and Management of Testosterone Deficiency: AUA Guideline. J Urol. Aug 2018; 200(2):423-432. PMID 29601923

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does not have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been developed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

Policy History/Revision	
Date	Description of Change
07/31/2026	Document became inactive.
01/01/2026	Document updated. The following changes were made to Coverage: 1) Removed reference to use in women; 2) removed NOTE 1; 3) Revised medically necessary criteria; 4) NOTE 2 moved to Policy guidelines; and 5) added (biological sex) to the experimental, investigational and/or unproven statement, intent unchanged. Added references 8 & 70.
02/01/2025	Document updated with literature change. Coverage unchanged. References 34, 49, 56, 65, and 69 added; others updated.
02/01/2024	Document updated with literature review. Coverage unchanged. References 13, 23, 24, 27-32, 34, 35, 37-40, 49, 65, 66 added; some removed; others revised.
01/01/2023	Reviewed. No changes.
05/01/2021	Document updated with literature review. The following change was made to Coverage: The experimental, investigational and/or unproven in all other situations statement was revised to include older men with type 2 diabetes mellitus and androgen deficiency. References 1, 8, 19-21, 23, 29, 50, 52 and 53 were added, others updated, and some removed.
03/15/2020	Reviewed. No changes.
08/01/2018	Document updated with literature review. The following change was made: NOTE 3 was replaced with EXCEPTION and the statement was revised to Individual consideration may be given for men continuously on therapy for at least one year and a morning total serum level or free testosterone level tested within the last year remains below or within the testing laboratory's normal range. References 10-11, 14, 22-24, 35-36, 40, 43-47, 57 added; some references removed.
04/15/2016	Reviewed. No changes.

12/15/2015	Document updated with literature review. The following was added in Coverage section: "NOTE 3: Individual consideration may be given for men currently on therapy when pretreatment levels are not available and the total serum level or free testosterone level remains below or within the testing laboratory's normal range." Otherwise, coverage unchanged.
02/12/2015	CPT/HCPCS code(s) updated. Testosterone therapy for women is considered experimental, investigational and/or unproven for ALL conditions, with the exception of treatment for metastatic breast cancer.
12/01/2014	Document updated with literature review. The following criteria was added to the medically necessary coverage statement: "Men requiring testosterone replacement therapy following post-bilateral-orchietomy" Otherwise, coverage remains unchanged.
10/01/2014	New medical document. Testosterone replacement therapy of one testosterone product may be considered medically necessary when the patient does not have any contraindications noted on the U.S. Food and Drug Administration-approved label for men with an established diagnosis of (primary or secondary) hypogonadism with androgen deficiency, when criteria is met; or HIV-infected men with low testosterone levels and hypogonadal symptoms or significant weight loss (e.g., > 5% lean body mass); or men on chronic-corticosteroid treatment with low testosterone levels and hypogonadal symptoms. Testosterone replacement therapy is considered experimental, investigational and/or unproven in all other situations in which the above criteria are not met, including but not limited to older men with low testosterone levels in the absence of clinical signs and symptoms of hypogonadism. CPT/HCPCS code(s) updated. This topic was previously addressed on RX501.007, Subcutaneous Hormone Implants.