

|                       |            |
|-----------------------|------------|
| Policy Number         | SUR703.005 |
| Policy Effective Date | 01/01/2026 |

## Heart Transplant

| Table of Contents                 |
|-----------------------------------|
| <a href="#">Coverage</a>          |
| <a href="#">Policy Guidelines</a> |
| <a href="#">Description</a>       |
| <a href="#">Rationale</a>         |
| <a href="#">Coding</a>            |
| <a href="#">References</a>        |
| <a href="#">Policy History</a>    |

| Related Policies (if applicable)                                                       |
|----------------------------------------------------------------------------------------|
| SUR703.006: Heart/Lung Transplant                                                      |
| SUR703.001: Organ and Tissue Transplantation (General Donor and Recipient Information) |
|                                                                                        |
|                                                                                        |
|                                                                                        |
|                                                                                        |

### Disclaimer

#### Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

### Legislative Mandates

**EXCEPTION: For Texas ONLY:** For policies (IFM, Student, Small Group, Mid-Market, Large Group, fully-insured Municipalities/Counties/Schools, State Employee Plans, PPO, HMO, POS) delivered, issued for delivery, or renewed on or after January 1, 2024, TIC Chapter 1380 (§§ 1380.001 – 1380.003 [SB 1040 Human Organ Transplant]) prohibits coverage of a human organ transplant or post-transplant care if the transplant operation is performed in China or another country known to have participated in forced organ harvesting; or the human organ to be transplanted was procured by a sale or donation originating in China or another country known to have participated in forced organ harvesting. The commissioner of state health services may designate countries who are known to have participated in forced organ harvesting. Forced organ harvesting is defined as the removal of one or more organs from a living person by means of coercion, abduction, deception, fraud, or abuse of power or a position of vulnerability.

### Coverage

Human heart transplant **may be considered medically necessary** in carefully selected individuals with end-stage heart failure who meet the United Network for Organ Sharing

(UNOS) guidelines for 1-6 Status and are not currently Inactive Status (formerly known as Status 7).

Heart retransplantation after a failed primary heart transplant **may be considered medically necessary** in individuals who meet criteria for heart transplantation.

Heart transplantation **is considered experimental, investigational and/or unproven** in all other situations.

## Policy Guidelines

### General Criteria

Potential contraindications for solid organ transplant subject to the judgment of the transplant center include the following:

- Known current malignancy, including metastatic cancer;
- Recent malignancy with a high risk of recurrence;
- Untreated systemic infection making immunosuppression unsafe, including chronic infection;
- Other irreversible end-stage diseases not attributed to heart or lung disease;
- History of cancer with a moderate risk of recurrence;
- Systemic disease that could be exacerbated by immunosuppression;
- Psychosocial conditions or chemical dependency affecting the ability to adhere to therapy.

Policy-specific potential contraindications include:

- Pulmonary hypertension that is fixed as evidenced by pulmonary vascular resistance >5 Wood units, or transpulmonary gradient  $\geq 16$  mm/Hg despite treatment<sup>a</sup>;
- Severe pulmonary disease, despite optimal medical therapy, not expected to improve with heart transplantation.<sup>a</sup>

<sup>a</sup> Some individuals may be candidates for combined heart and lung transplantation (see medical policy SUR703.006).

### Cardiac-Specific Criteria

Specific criteria for prioritizing donor thoracic organs for transplant are provided by the Organ Procurement and Transplantation Network (OPTN) and implemented through a contract with UNOS. Donor thoracic organs are prioritized by UNOS on the basis of recipient medical urgency, distance from donor hospital, and pediatric status. Individuals who are most severely ill (status 1A) are given the highest priority. The following factors are considered in assessing the severity of illness: reliance on continuous mechanical ventilation, infusion of intravenous inotropes, and/or dependency on mechanical circulatory support (ie, total artificial heart, intra-aortic balloon pump, extracorporeal membrane oxygenator, ventricular assist device).

Additional criteria, which are considered in pediatric individuals, include diagnosis of an OPTN-approved congenital heart disease, presence of ductal dependent pulmonary or systemic

circulation, and diagnosis of hypertrophic or restrictive cardiomyopathy while less than 1-year-old. Of note, pediatric heart transplant candidates who remain on the waiting list at the time of their 18th birthday without receiving a transplant continue to qualify for medical urgency status based on the pediatric criteria.

Specific criteria for prioritizing donor thoracic organs for retransplant include severe coronary allograft vasculopathy, mild or moderate coronary allograft vasculopathy with a left ventricular ejection fraction less than 45%, coronary allograft vasculopathy with restrictive physiology, or symptomatic graft dysfunction without evidence of active rejection.

## Description

A heart transplant and a retransplant consist of replacing a diseased heart with a healthy donor heart. Transplantation is used for individuals with refractory end-stage cardiac disease.

### **Solid Organ Transplantation**

Solid organ transplantation offers a treatment option for patients with different types of end-stage organ failure that can be lifesaving or provide significant improvements to a patient's quality of life. (1) Many advances have been made in the last several decades to reduce perioperative complications. Available data support improvement in long-term survival as well as improved quality of life, particularly for liver, kidney, pancreas, heart, and lung transplants. Allograft rejection remains a key early and late complication risk for any organ transplantation. Transplant recipients require life-long immunosuppression to prevent rejection. Patients are prioritized for transplant by mortality risk and severity of illness criteria developed by the Organ Procurement and Transplantation Network and United Network for Organ Sharing (UNOS).

### Heart Transplant

In 2024, 48,136 transplants were performed in the United States procured from over 41,000 deceased donors and over 7,000 living donors. (2) Heart transplants were the third most common procedure with 4,560 transplants performed from both deceased and living donors in 2024. As of July 2025, there were 3,822 patients on the waiting list for a heart transplant. (3) Rose et al. (2024) reported a 62% lower rate of heart transplants among women compared with men and a 46% lower rate in Black men compared with White men in a retrospective database review from 2010 to 2018. (4)

Most heart transplant recipients now are hospitalized as status 1 patients at the time of transplant. This shift has occurred due to the increasing demand for the scarce resource of donor organs resulting in an increased waiting time for recipients. Patients initially listed as status 2 candidates may deteriorate to a status 1 candidate before a donor organ becomes available. Alternatively, as medical and device therapy for advanced heart failure improves, some patients on the transplant list will recover enough function to be delisted. Lietz and Miller (2007) reported on survival for patients on the heart transplant waiting list, comparing the era between 1990 and 1994 with the era of 2000 to 2005. (5) One-year survival for a UNOS status 1

candidate improved from 49.5% to 69.0%. Status 2 candidates fared even better, with 89.4% surviving 1 year compared with 81.8% in the earlier time period.

Johnson et al. (2010) reported on waiting list trends in the U.S. between 1999 and 2008. (6) The proportion of patients listed as status 1 increased, even as the waiting list and posttransplant mortality for this group have decreased. Meanwhile, status 2 patients have decreased as a proportion of all candidates. Completed transplants have trended toward the extremes of age, with more infants and patients older than age 65 years having transplants in recent years. Bakhtiyar et al (2020) evaluated survival among patients (N=95,323) wait-listed for heart transplantation between January 1, 1987 and December 29, 2017 using UNOS data. (7) Results revealed 1-year survival on the wait list increased from 34.1% in 1987 to 1990 to 67.8% in 2011 to 2017 (difference in proportions, 0.34%; 95% confidence interval [CI], 0.32% to 0.36%;  $p<.001$ ). One-year wait list survival also significantly increased for candidates with ventricular assist devices from 10.2% in 1996 to 2000 to 70% in 2011 to 2017 (difference in proportions, 0.60%; 95% CI, 0.58% to 0.62%;  $p<.001$ ).

Alshawabkeh et al. (2018) reported on the 1-year probability of the combined outcome of death or delisting due to clinical worsening for patients on the heart transplant waiting list, comparing the periods of April 1, 1986 to January 19, 1999 (early era) and January 20, 1999 to June 2, 2014 (current era). (8) For adults without congenital heart disease (CHD), the probability of the combined outcome was lower in the current era compared with the early era, regardless of whether the patient was listed in status 1 (14.5% vs. 22.7%;  $p<.0001$ ) or 2 (9.0% vs. 12.8%,  $p<.0001$ ). When comparing the current and early eras in adults with CHD, a reduction in the probability of the combined outcome was demonstrated in those listed in status 1 (17.6% vs. 43.3%, respectively;  $p<.0001$ ), whereas the outcome remained unchanged for those listed in status 2 (10.6% vs. 10.4%, respectively;  $p=.94$ ).

In adults with CHD, factors associated with waitlist death or delisting due to clinical worsening within 1 year were also examined by Alshawabkeh et al. (2016). (9) A multivariate analysis identified that an estimated glomerular filtration rate less than 60 ml/min/1.73 m<sup>2</sup> (hazard ratio [HR], 1.4; 95% CI, 1.0 to 1.9;  $p=.043$ ), albumin less than 3.2 g/dl (HR, 2.0; 95% CI, 1.3 to 2.9;  $p<.001$ ), and hospitalization at the time of listing in the intensive care unit (HR, 2.3; 95% CI, 1.6 to 3.5;  $p<.001$ ) or a non-intensive care hospital unit (HR, 1.9; 95% CI, 1.2 to 3.0;  $p=.006$ ) were associated with waitlist death or delisting due to clinical worsening within 1 year.

Magnetta et al. (2019) reported outcomes for children on the heart transplant waiting list, comparing the periods of December 16, 2011 to March 21, 2016 (era 1) and March 22, 2016 to June 30, 2018 (era 2). (10) There was a significant decrease from era 1 to era 2 in the proportion of patients listed as status 1 (70% vs. 56%;  $p<.001$ ), while the proportion of patients with CHD significantly increased across eras (49% to 54%;  $p=.018$ ). The median time on the waitlist increased from 68 days to 78 days ( $p=.005$ ). There were no significant differences across eras in the cumulative incidence of death on the waitlist among all candidates (subdistribution HR, 0.96; 95% CI, 0.80 to 1.14;  $p=.63$ ) and among those listed status 1A (subdistribution HR,

1.16; 95% CI, 0.95 to 1.41;  $p=.14$ ). Graft survival at 90 days was also similar across eras in the overall population and in those with CHD ( $p>.53$  for both).

As a consequence, aggressive treatment of heart failure has been emphasized in recent guidelines. Prognostic criteria have been investigated to identify patients who have truly exhausted medical therapy and thus are likely to derive the maximum benefit for heart transplantation. Maximal oxygen consumption ( $Vo_2max$ ), which is measured during maximal exercise, is a measure suggested as a critical objective criterion of the functional reserve of the heart. The American College of Cardiology and American Heart Association have adopted  $Vo_2max$  as a criterion for patient selection. (11)

Methods other than  $Vo_2max$  have been proposed as predictive models in adults. (12-15) The Heart Failure Survival Scale and the Seattle Heart Failure Model (SHFM) are examples. In particular, the SHFM provides an estimate of 1-, 2-, and 3-year survival with the use of routinely obtained clinical and laboratory data. Information on pharmacologic and device usage is incorporated into the model, permitting some estimation on the effects of current, more aggressive heart failure treatment strategies. Levy et al. (2006) introduced the model using a multivariate analysis of data from the Prospective Randomized Amlodipine Survival Evaluation-1 heart failure trial ( $N=1125$ ). (16) Applied to the data of 5 other heart failure trials, SHFM correlated well with actual survival ( $r=0.98$ ). SHFM has been validated in both ambulatory and hospitalized heart failure populations, (17-19) but with a noted underestimation of mortality risk, particularly in Black adults and device recipients. (20, 21) None of these models has been universally adopted by transplant centers.

### **Regulatory Status**

Solid organ transplants are a surgical procedure and, as such, are not subject to regulation by the U.S. Food and Drug Administration (FDA).

The FDA regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation Title 21, parts 1270 and 1271. Solid organs used for transplantation are subject to these regulations.

## **Rationale**

Medical policies assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Due to the nature of the population discussed herein, there are no RCTs comparing heart transplantation with alternatives, including left ventricular assist devices (LVADs). Systematic reviews are based on case series and registry data. Randomized controlled trials have been published on related topics (e.g., comparing surgical technique, infection prophylaxis regimens, or immunosuppressive therapy) but are not germane to this medical policy.

### **Initial Heart Transplant**

#### **Clinical Context and Therapy Purpose**

In the U.S., approximately 6 million people 20 years of age and older have heart failure and 1 in 8 deaths have heart failure mentioned on the death certificate. (22) The reduction of cardiac output is considered to be severe when systemic circulation cannot meet the body's needs under minimal exertion.

Heart failure may be due to a number of differing etiologies, including ischemic heart disease, cardiomyopathy, or congenital heart disease (CHD). The leading indication for a heart transplant has shifted over time from ischemic to nonischemic cardiomyopathy. From 2009 to 2014, nonischemic cardiomyopathy was the dominant underlying primary diagnosis among recipients 18 to 39 years (64%) and 40 to 59 years (51%) undergoing transplant operations. (23) Ischemic cardiomyopathy was the dominant underlying primary diagnosis among heart transplant recipients 60 to 69 years (50%) and 70 years and older (55%). Overall, ischemic cardiomyopathy is the underlying heart failure diagnosis in approximately 40% of men and 20% of women who receive a transplant. Approximately 3% of heart transplants during this time period were in adults with CHD.

The purpose of a heart transplant in individuals who have end-stage heart failure is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

#### ***Populations***

The relevant population of interest is individuals who have end-stage heart failure.

### *Interventions*

The therapy being considered is a heart transplant.

### *Comparators*

The following therapies and practices are currently being used to make decisions about reducing the risk of end-stage heart failure: guideline-directed medical therapy; surgery including coronary bypass surgery, heart valve repair or replacement, and ventricular assist devices.

### *Outcomes*

The general outcomes of interest are overall survival (OS), symptoms, and morbid events (e.g., immunosuppression, graft failure, surgical complications, infections, cardiovascular complications, malignancies). See the Potential Contraindications section for a detailed discussion of outcomes in individuals with malignancy, HIV, older age, pulmonary hypertension, and renal insufficiency, and children with intellectual disability. Follow-up of 1, 2, 5, and 10 years is of interest for heart transplant outcomes.

### Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

### Retrospective Studies

A study by Jaramillo et al. (2013) examined characteristics of patients who survived more than 20 years after heart transplantation at a single-center in Spain. (24) Thirty-nine heart transplant recipients who survived over 20 years posttransplant were compared with 98 patients who died between 1 and 20 years posttransplant. Independent factors associated with long-term survival were younger recipient age (i.e., <45 years vs. ≥45 years; odds ratio [OR], 3.9; 95% confidence interval [CI], 1.6 to 9.7) and idiopathic cardiomyopathy (i.e., vs. other etiologies; OR, 3.3; 95% CI, 1.4 to 7.8).

### Registry Studies

According to the Organ Procurement and Transplantation Network (OPTN), 1-year Kaplan-Meier survival estimates for heart transplants performed between 2008 and 2015, based on available U.S. data as of June 14, 2024 were 90.3% (95% CI, 89.6% to 90.9% ) for men and 90.7% (95% CI, 89.6% to 91.7% ) for women. (3) The 3-year survival rates were 84.7% (95% CI, 83.8% to 85.5%) for men and 84.1% (95% CI, 82.7% to 85.4%) for women, and the 5-year survival rates were 77.8% (95% CI, 76.8% to 78.8%) and 75.9% (95% CI, 74.2% to 77.6%), respectively. There

was no major difference in 1-, 3-, and 5-year survival rates between different age groups among adult recipients (Table 1).

**Table 1. Kaplan-Meier Patient Survival Rates for Heart Transplants Performed From 2008 to 2015**

| Recipient Age | Years Posttransplant |                           |                      |                           |                      |                           |
|---------------|----------------------|---------------------------|----------------------|---------------------------|----------------------|---------------------------|
|               | 1 Year <sup>a</sup>  |                           | 3 Years <sup>a</sup> |                           | 5 Years <sup>a</sup> |                           |
|               | No. Alive            | Survival Rate (95% CI), % | No. Alive            | Survival Rate (95% CI), % | No. Alive            | Survival Rate (95% CI), % |
| <1 year       | 406                  | 87.4 (84.0 to 90.1)       | 362                  | 83.9 (80.1 to 87.0)       | 317                  | 75.0 (70.6 to 78.8)       |
| 1-5 years     | 345                  | 92.4 (89.2 to 94.6)       | 282                  | 85.0 (80.7 to 88.4)       | 257                  | 78.1 (73.3 to 82.1)       |
| 6-10 years    | 223                  | 91.9 (87.7 to 94.7)       | 186                  | 87.0 (81.7 to 90.8)       | 166                  | 84.1 (78.3 to 88.5)       |
| 11-17 years   | 507                  | 96.6 (94.7 to 97.9)       | 458                  | 91.5 (88.7 to 93.6)       | 356                  | 78.4 (74.4 to 81.9)       |
| 18-34 years   | 840                  | 90.9 (88.8 to 92.6)       | 722                  | 82.1 (79.4 to 84.5)       | 597                  | 72.3 (69.2 to 75.2)       |
| 35-49 years   | 1588                 | 90.6 (89.1 to 91.8)       | 1399                 | 84.8 (83.0 to 86.4)       | 1233                 | 78.2 (76.1 to 80.1)       |
| 50-64 years   | 3896                 | 90.4 (89.5 to 91.3)       | 3373                 | 84.8 (83.7 to 85.9)       | 2972                 | 78.3 (76.9 to 79.5)       |
| 65+ years     | 1514                 | 88.2 (86.6 to 89.6)       | 1191                 | 82.1 (80.0 to 83.9)       | 879                  | 75.4 (72.9 to 77.8)       |

Source: Organ Procurement and Transplantation Network. (3)

CI: confidence interval; No: number; NR: not reported.

<sup>a</sup> One-year survival based on 2012-2015 transplants, 3-year survival based on 2010-2013 transplants, 5-year survival based on 2008-2011 transplants.

Nguyen et al. (2017) investigated the benefit of heart transplantation compared with surveillance while on a waiting list while accounting for the estimated risk of a given donor-recipient match among 28,548 heart transplant candidates in OPTN between 2006 and 2015. (25) The net benefit from heart transplantation was evident across all estimates of donor-recipient status 1A candidates (lowest risk quartile hazard ratio [HR], 0.37; 95% CI, 0.31 to 0.43; highest-risk quartile HR, 0.52; 95% CI, 0.44 to 0.61) and status 1B candidates (lowest-risk quartile HR, 0.41; 95% CI, 0.36 to 0.47; highest-risk quartile HR, 0.66; 95% CI, 0.58 to 0.74). Status 2 candidates also showed a benefit from heart transplantation; however, the survival benefit was delayed. For the highest-risk donor-recipient matches, a net benefit of transplantation occurred immediately for status 1A candidates, after 12 months for status 1B candidates, and after 3 years for status 2 candidates.

Rana et al. (2015) retrospectively analyzed solid organ transplant recipients registered in the United Network for Organ Sharing (UNOS) database from 1987 to 2012, including 54,746 patients who underwent a heart transplant. (26) Transplant recipients were compared with patients listed for transplant but who did not receive one; heart recipients were awarded the transplant based on propensity score matching, which served to measure a variety of clinical characteristics. After matching, the median survival was 9.5 years in transplant recipients compared with 2.1 years in waiting list patients.

Several studies have analyzed factors associated with survival in heart transplant patients. For example, Lund et al. (2016) examined the risk factors associated with 10-year posttransplant mortality among patients undergoing heart transplantation between 2000 and 2005 using the International Society for Heart and Lung Transplantation (ISHLT) Registry. (23) Markers of pretransplant severity of illness, such as pretransplant ventilator use (HR, 1.35; 95% CI, 1.17 to 1.56; n=338), dialysis use (HR, 1.51; 95% CI, 1.28 to 1.78; n=332), underlying diagnoses of ischemic (HR, 1.16; 95% CI: 1.10 to 1.23; n=7822), congenital (HR, 1.21; 95% CI, 1.04 to 1.42; n=456) or restrictive (HR, 1.33; 95% CI, 1.13 to 1.58; n=315) heart disease (vs. nonischemic cardiomyopathy), and retransplant (HR, 1.18; 95% CI, 1.02 to 1.35; n=489) were associated with posttransplant mortality risk at 10 years.

A study by Kilic et al. (2012) analyzed prospectively collected data from the UNOS registry. (27) The analysis included 9404 patients who had survived 10 years after a heart transplant and 10,373 patients who had died before 10 years. Among individuals who had died, the mean survival was 3.7 years posttransplant. In multivariate analysis, statistically significant predictors of surviving at least 10 years after heart transplant included age younger than 55 years (OR, 1.24; 95% CI, 1.10 to 1.38), younger donor age (OR, 1.01; 95% CI, 1.01 to 1.02), shorter ischemic time (OR, 1.11; 95% CI, 1.05 to 1.18), White race (OR, 1.35; 95% CI, 1.17 to 1.56), and annual center volume of 9 or more heart transplants (OR, 1.31; 95% CI, 1.17 to 1.47). Factors that significantly decreased the likelihood of 10-year survival in multivariate analysis included the use of mechanical ventilation (OR, 0.53; 95% CI, 0.36 to 0.78) and diabetes (OR, 0.67; 95% CI, 0.57 to 0.78).

## Pediatric Considerations

### *Retrospective Studies*

An analysis of data from the Pediatric Heart Transplant Study (2013), which includes data on all pediatric transplants at 35 participating institutions, suggests that 5-year survival for pediatric heart transplants has improved over time (76% for patients transplanted from 2000 to 2004 vs. 83% for patients transplanted from 2005 to 2009). (28)

Auerbach et al. (2012) published a retrospective review of pediatric cardiac transplantation patients. (29) A total of 191 patients who underwent primary heart transplantation at a single-center in the U.S. were included; their mean age was 9.7 years (range, 0 to 23.6 years). Overall graft survival was 82% at 1 year and 68% at 5 years; the most common causes of graft loss were acute rejection and graft vasculopathy. Overall survival was 82% at 1 year and 72% at 5 years. In multivariate analysis, the authors found that CHD (HR, 1.6; 95% CI, 1.02 to 2.64) and mechanical

ventilation at the time of transplantation (HR, 1.6; 95% CI, 1.13 to 3.10) were both significantly and independently associated with an increased risk of graft loss. Renal dysfunction was a significant risk factor in univariate analysis but was not included in the multivariate model due to the small size of the study group. Study limitations included the retrospective design and single-center sample.

### *Registry Studies*

According to OPTN, patients between the ages of 11 and 17 years of age had the highest 1- and 3-year survival rates among pediatric patients who underwent a heart transplant in the U.S. between 2008 and 2015. (3) Patients younger than 1 year of age had the lowest 1-, 3-, and 5-year survival rates among pediatric patients (Table 1).

Rossano et al. (2016) examined survival among pediatric heart transplant recipients using the ISHLT Registry. (30) Among 12,091 pediatric patients undergoing heart transplantation between 1982 and 2014, the overall median survival was 20.7 years for infants (n=2994), 18.2 years for children between the ages of 1 to 5 years (n=2720), 14.0 years for those 6 to 10 years (n=1743), and 12.7 years for those 11 to 17 years (n=4684). Because the first year posttransplant represents the greatest risk for mortality, survival conditional on survival to 1 year was longer.

Rossano et al conducted a multivariable analysis of pediatric patients undergoing a heart transplant between 2003 and 2013 to identify the factors associated with 1-year mortality. (30) Infection requiring intravenous drug therapy within 2 weeks of transplant (HR, 1.36; 95% CI, 1.10 to 1.68; n=681), ventilator use (HR, 1.41; 95% CI, 1.13 to 1.76; n=826), donor cause of death (cerebrovascular accident vs. head trauma; HR, 1.59; 95% CI, 1.20 to 2.09; n=396), diagnosis (CHD vs. cardiomyopathy; HR, 1.91; 95% CI, 1.46 to 2.52; n=1979; retransplant vs. cardiomyopathy; HR, 2.23; 95% CI, 1.53 to 3.25; n=304), recipient dialysis (HR, 2.36; 95% CI, 1.57 to 3.57; n=146), extracorporeal membrane oxygenation (ECMO) with a diagnosis of CHD versus no ECMO (HR, 2.42; 95% CI, 1.74 to 3.35; n=145), ischemic time (p<.001), donor weight (p<.001), estimated glomerular filtration rate (eGFR; p=.002), and pediatric center volume (p<.001) were risk factors for 1-year mortality. Earlier era (1999 to 2000 vs. 2007 to 2009), CHD (vs. dilated cardiomyopathy), use of ECMO (vs. no device), and pediatric center volume were risk factors for 5-, 10-, and 15-year mortality. A panel-reactive antibody greater than 10% was associated with worse 5- and 10-year survival and eGFR was associated with 5- and 10-year mortality.

A retrospective analysis of the OPTN data focusing on the adolescent population was reported by Savla et al. (2014). (31) From 1987 to 2011, heart transplants were performed in 99 adolescents (age, 13 to 18 years) with myocarditis and 456 adolescents with CHD. Among transplant recipients with myocarditis, median graft survival was 6.9 years (95% CI, 5.6 to 9.6 years), which was significantly lower than other age groups (ie, 11.8 years and 12.0 years in younger and older adults, respectively). However, adolescents with CHD had a graft survival rate of 7.4 years (95% CI, 6.8 to 8.6 years), similar to that of other age groups.

Noting that children listed for heart transplantation have the highest waiting list mortality of all solid organ transplant patients, Almond et al (2009) analyzed data from the U.S. Scientific Registry of Transplant Recipients to determine whether the pediatric heart allocation system, as revised in 1999, was prioritizing patients optimally and to identify high-risk populations that may benefit from pediatric cardiac assist devices. (32) Of 3098 children (<18 years of age) listed between 1999 and 2006, 1874 (60%) were listed as status 1A. Of these 1874, 30% were placed on ventilation, and 18% were receiving ECMO. Overall, 533 (17%) died, 1943 (63%) received transplants, 252 (8%) recovered, and 370 (12%) remained listed. The authors found that status 1A patients were a heterogeneous population with a large variation in mortality based on patient-specific factors. Predictors of waiting list mortality included ECMO support (HR, 3.1), ventilator support (HR, 1.9), listing status 1A (HR, 2.2), CHD (HR, 2.2), dialysis support (HR, 1.9), and non-White race/ethnicity (HR, 1.7). The authors concluded that the pediatric heart allocation system was capturing medical urgency poorly, specific high-risk subgroups could be identified, and further research would be needed to better define the optimal organ allocation system for pediatric heart transplantation.

#### Section Summary: Initial Heart Transplant

The evidence supports a net benefit for heart transplantation compared with waitlist for status 1A and 1B candidates. Status 2 candidates also show a benefit from heart transplantation; however, the survival benefit is delayed. Data from national and international registries have found high patient survival rates after initial heart transplant among adult and pediatric patients (e.g., a 5-year survival rate, 78%).

### **Heart Retransplantation**

#### Clinical Context and Therapy Purpose

From 2008 to 2015, approximately 4% of heart transplants were repeated transplantations. (3) Heart retransplantation raises ethical issues due to the lack of sufficient donor hearts for initial transplants. The UNOS does not have separate organ allocation criteria for repeat heart transplant recipients.

The purpose of heart retransplants in individuals who have had a prior heart transplant complicated by graft failure or severe heart dysfunction is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

#### *Populations*

The relevant population of interest is individuals who have had a prior heart transplant complicated by graft failure or severe heart dysfunction.

#### *Interventions*

The therapy being considered is a heart retransplant.

#### *Comparators*

The following therapies and practices are currently being used to make decisions about reducing the risk of end-stage heart failure: guideline-directed medical therapy; surgery including coronary bypass surgery, heart valve repair or replacement, and ventricular assist devices.

### *Outcomes*

The general outcomes of interest are OS, symptoms, and morbid events (e.g., immunosuppression, graft failure, surgical complications, infections, cardiovascular complications, malignancies). See the Potential Contraindications section for a detailed discussion of outcomes in individuals with malignancy, HIV, older age, pulmonary hypertension, and renal insufficiency, and children with intellectual disability. Follow-up of 1, 2, 5, and 10 years is of interest for heart transplant outcomes.

### Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

### Systematic Reviews

A number of studies have reviewed the clinical experience with heart retransplantation in adults. Tjang et al. (2008) published a systematic review of the literature on the clinical experience with adult heart retransplantation; reviewers identified 22 studies. (33) The most common indications for retransplantation were cardiac allograft vasculopathy (55%), acute rejection (19%), and primary graft failure (17%). The early mortality rate in individual studies was 16% (range, 5% to 38%). Some factors associated with poorer outcomes after retransplantation were shorter transplant interval, refractory acute rejection, primary graft failure, and an initial diagnosis of ischemic cardiomyopathy.

### Retrospective Reviews

Zhu et al. (2022) evaluated outcomes after heart retransplantation for 123 patients (112 adult and 11 pediatric patients) as compared to those who received a primary heart transplant at a single-center over a 50-year period (January 6, 1968 to June 2019). (34) The indications for retransplantation included cardiac allograft vasculopathy (80%), primary graft dysfunction (15%), and refractory acute rejection (5%). The mean time interval between the primary and retransplant was 6.4 years. Patients who underwent a retransplantation were significantly more likely to have hypertension (73.3% vs. 53.3%;  $p=.0022$ ), hyperlipidemia (66.7% vs. 30.7%;  $p<.0001$ ), and require dialysis (11.7% vs. 2.9%;  $p=.0025$ ) as compared to those undergoing a primary heart transplant. After matching, postoperative outcomes and complications including hospital stay (mean 22.9 vs. 25.8 days;  $p=.49$ ), intensive care unit stay (mean 12.2 vs. 9.9 days;

p=.48), respiratory failure (41.7% vs. 20.6%; p=.083), dialysis (21.2% vs. 24.2%; p=.82), pneumonia (12.9% vs. 9.6%; p=.48), septicemia (1.6% vs. 9.4%; p=.10), and rejection within the first year after transplantation requiring hospitalization (21.5% vs. 26.2%; p=.82) were similar between the retransplant and primary transplant groups, respectively. Matched median survival after retransplantation was 4.6 years versus 6.5 years after primary heart transplantation (p=.36).

In a retrospective review, Saito et al. (2013) evaluated 593 patients with heart transplants performed at their institution, 22 (4%) of whom required retransplants. (35) The mean interval between initial and repeat transplants was 5.1 years. The indications for a repeat transplant were acute rejection in 7 (32%) patients, graft vascular disease in 10 (45%) patients, and primary graft failure in 5 (23%) patients. The 30-day mortality rate after cardiac retransplantation was 32% (7/22 patients). Among patients who survived the first 30 days (n=15), 1-, 5-, and 10-year survival rates were 93.3%, 79%, and 59%, respectively. Comparable survival rates for patients undergoing primary cardiac transplants at the same institution (n=448) were 93%, 82%, and 63%, respectively. An interval of 1 year or less between the primary and repeat transplantation significantly increased the risk of mortality. Three of 9 (33.3%) patients with less than 1 year between the primary and retransplantation survived to 30 days; by comparison, 12 (92%) of 13 patients with at least 1 year between primary and retransplantation were alive at 30 days after surgery.

#### Registry Studies

An analysis of OPTN data from 2008 to 2015 found that 724 (3.9%) retransplants (of 18,676 heart transplants) were performed. Kaplan-Meier patient survival estimates at 1, 3, and 5 years were lower among the retransplant recipients than among primary transplant recipients (Table 2).

**Table 2. Kaplan-Meier Patient Survival Estimates for Primary and Repeat Heart Transplants Performed Between 2008 and 2015**

| Years Posttransplant | Transplant Type    |                               |              |                   |                               |              |
|----------------------|--------------------|-------------------------------|--------------|-------------------|-------------------------------|--------------|
|                      | Primary Transplant |                               |              | Repeat Transplant |                               |              |
|                      | No. Alive          | Survival Rate, % <sup>a</sup> | 95% CI, %    | No. Alive         | Survival Rate, % <sup>a</sup> | 95% CI, %    |
| 1 year               | 9013               | 90.9                          | 90.3 to 91.4 | 320               | 87.0                          | 83.1 to 90.0 |
| 3 years              | 7711               | 85.6                          | 84.8 to 86.3 | 286               | 76.5                          | 71.8 to 80.4 |
| 5 years              | 6572               | 78.6                          | 77.7 to 79.4 | 237               | 69.7                          | 64.6 to 74.2 |

Source: Organ Procurement and Transplantation Network. (3)

CI: confidence interval; No: number.

<sup>a</sup> One-year survival rates based on 2012-2015 transplants, 3-year survival rates based on 2010-2013 transplants, 5- year survival rates based on 2008-2011 transplants.

In a study analyzing UNOS data from January 1996 and November 2017, Miller et al. (2019) reported that 349 (0.6%) early/acute retransplants (occurring ≤1 year after the previous

transplant) and 2202 (3.5%) late retransplants (occurring >1 year after the previous transplant) were performed from a sample of 62,112 heart transplants. (36) Compared with a matched group of patients undergoing initial transplantation, patients undergoing late retransplantation were not at an increased risk of death (HR, 1.08;  $p=.084$ ) or the combined outcome of death or retransplantation (HR, 1.07;  $p=.114$ ). Additionally, patients undergoing late retransplant had comparable rates of 1-year all-cause mortality when compared to patients undergoing initial transplant (13.8% vs. 14.5%, respectively;  $p=.517$ ). Conversely, patients undergoing early/acute transplant had higher rates of 1-year all-cause mortality when compared to patients undergoing initial transplant (35% vs. 21.6%;  $p<.001$ ). Furthermore, early/acute retransplantation was associated with an increased risk of all-cause mortality (HR, 1.79;  $p<.001$ ) and the combined outcome of death or retransplantation (HR, 1.72;  $p<.001$ ).

Goldraich et al. (2016) examined the survival data for adult heart recipients with cardiac allograft vasculopathy who were retransplanted ( $n=65$ ) or managed medically ( $n=4530$ ). (37) During a median follow-up of 4 years, 24 deaths occurred among those who underwent retransplantation and 1466 deaths among those medically managed. There was no significant difference in survival rates at 9 years (55% in retransplant recipients vs. 51% in medically managed patients,  $p=.88$ ). In a subgroup analysis, the retransplant group ( $n=65$ ) had longer survival than the medically managed group at 1 year after the development of coronary allograft vasculopathy ( $n=124$ ;  $p=.02$ ).

In an analysis of the OPTN data from 1995 to 2012, Belli et al. (2014) reported that 987 (3.5%) retransplants were performed from a sample of 28,464 heart transplants. (38) Median survival among retransplant recipients was 8 years. The estimated survival rates at 1, 5, 10, and 15 years following retransplant were 80%, 64%, 47%, and 30%, respectively. Compared with primary transplant recipients, retransplant patients had a somewhat higher risk of death (relative risk, 1.27, 95% CI, 1.13 to 1.42).

In a study analyzing UNOS data, Friedland-Little et al. (2014) reported no survival differences between third and second transplants (76% for third transplant vs. 80% for second transplant at 1 year; 62% for third transplant vs. 58% for second transplant at 5 years; 53% for third transplant vs. 34% for second transplant at 10 years,  $p=.73$ ). (39) However, study conclusions might have been limited because of the small number ( $n=25$ ) of third heart transplants.

### Pediatric Considerations

As with initial heart transplants, children awaiting heart retransplantation have high waitlist mortality. A study by Bock et al. (2015) evaluated data on 632 pediatric patients who were listed for a heart retransplant for at least 1 year (median, 7.3 years) after the primary transplant. (40) Patients' median age was 4 years at the time of the primary transplant and 14 years when relisted. Median waiting time was 75.3 days, and the mortality rate was 25.2% (159/632). However, waitlist mortality decreased significantly after 2006 (31% before 2006 and 17% after 2006,  $p<.01$ ).

Conway et al. (2014) analyzed the ISHLT Registry to compare the outcomes after retransplantation with primary heart transplantation among pediatric (<18 years of age) transplant recipients from 1998 to 2010. (41) Of the 9882 heart transplant recipients with available clinical outcomes data, 9248 (93.6%) were primary transplants, 602 (6.1%) were retransplants (second graft), and 32 (0.3%) were third or fourth grafts. The median ages at primary transplant and retransplant were 7 years (range, 0 to 14 years) and 14 years (range, 1 to 26 years), respectively. The mean intertransplant interval was 6.8 years after primary transplant. The most common indications for retransplantation were coronary allograft vasculopathy (n=352 [59%]), nonspecific graft failure (n=52 [9%]), and acute rejection (n=49 [8%]). Retransplantation was associated with similar early survival but decreased long-term survival compared with initial transplantation. After primary transplantation, the survival rate was 84% at 1 year, 72% at 5 years, 60% at 10 years, and 42% at 20 years, compared with 81% at 1 year, 63% at 5 years, 46% at 10 years, and 26% at 20 years after retransplantation, respectively. The median survival rate was longer in primary transplant recipients, reaching 15 years (vs. 8.7 years after retransplantation). The most common causes of death after retransplantation were cardiovascular other than vasculopathy (28%), graft failure (10%), infection (9%), noncardiac organ failure (9%), coronary allograft vasculopathy (4%), and acute rejection (3%).

#### Section Summary: Heart Retransplantation

In both adult and pediatric studies, poorer survival after retransplantation compared with initial transplantation is not surprising given that patients undergoing retransplantation experienced additional clinical disease or adverse events. Data from national and international registries have found high patient survival rates after heart retransplant among adult and pediatric patients (e.g., a 5-year survival rate, 69%). Cardiac allograft vasculopathy is the most common indication for heart retransplantation among adult and pediatric patients. Considering the scarcity of heart donors and the few treatment options for cardiac allograft vasculopathy, additional studies must be done to further examine the survival benefit of cardiac retransplantation over medical management among patients with cardiac allograft vasculopathy.

#### **Potential Contraindications to Heart Transplant/Retransplantation**

Individual transplant centers may differ in their guidelines, and individual patient characteristics may vary within a specific condition. In general, heart transplantation is contraindicated in patients who are not expected to survive the procedure or in whom patient-oriented outcomes (e.g., morbidity, mortality) are not expected to change due to comorbid conditions unaffected by transplantation (e.g., imminently terminal cancer, another disease). Moreover, consideration is given to conditions in which the necessary immunosuppression would lead to hastened demise, such as active untreated infection. However, stable chronic infections have not always been shown to reduce life expectancy in heart transplant patients.

#### Malignancy

Pretransplant malignancy is considered a relative contraindication for heart transplantation because malignancy has the potential to reduce life expectancy and could prohibit immune

suppression after transplantation. However, with improved cancer survival and use of cardiotoxic chemotherapy and radiotherapy, the need for heart transplantation has increased in this population, Mistiaen (2015) conducted a systematic review to study posttransplant outcomes of patients with pretransplant malignancy. (42) Most selected studies were small case series (median sample size, 17 patients; range, 7 to 1117 patients; mean age range, 6 to 52 years). Hematologic malignancy and breast cancer were the most common types of pretransplant malignancies. Dilated, congestive, or idiopathic cardiomyopathy were the most common reasons for transplantation in 4 case series, chemotherapy-related cardiomyopathy was the most important reason for transplantation in the other series. Hospital mortality rates varied between 0% and 33%, with small sample size potentially explaining the observed variation.

Yoosabai et al. (2015) retrospectively reviewed data on 23,171 heart transplant recipients in the OPTN/UNOS database to identify whether pretransplant malignancy increases the risk of post-transplant malignancy. (43) Posttransplant malignancy was diagnosed in 2673 (11.5%) recipients during the study period. A history of any pretransplant malignancy was associated with an increased risk of overall post-transplant malignancy (subhazard ratio, 1.51;  $p<.01$ ), skin malignancies (subhazard ratio, 1.55,  $p<.01$ ), and solid organ malignancies (subhazard ratio, 1.54,  $p<.01$ ) on multivariate analysis.

One large series by Oliveira et al. (2012) reported similar short- and long-term post-transplant survival rates for patients who received chemotherapy-related ( $n=232$ ) and for those with another nonischemic-related cardiomyopathy ( $n=8890$ ). (44) The 1-, 3-, and 5-year survival rates were 86%, 79%, and 71% for patients with chemotherapy-related cardiomyopathy compared with 87%, 81%, and 74% for other transplant patients, respectively. Similar 1-year survival findings were observed in smaller series. Two-, 5-, and 10-year survival rates among patients with pretransplant malignancy were also comparable with other transplant patients. In addition to the non-malignancy-related factors such as cardiac, pulmonary, and renal dysfunction, 2 malignancy-related factors were identified as independent predictors of 5-year survival. Malignancy-free interval (the interval between treatment of cancer and heart transplantation) of less than 1 year was associated with a lower 5-year survival rate ( $<60\%$ ) than with a longer interval ( $>75\%$ ). Patients with prior hematologic malignancies had increased posttransplant mortality rates in 3 small series. Recurrence of malignancy was more frequent among patients with a shorter disease-free interval (63%, 26%, and 6% among patients with  $<1$  year, 1 to 5 years, and  $>5$  years of disease-free interval, respectively). (45)

The evaluation of a candidate who has a history of cancer must consider the prognosis and risk of recurrence from available information including tumor type and stage, response to therapy, and time since therapy was completed. Although evidence is limited, patients for whom cancer is thought to be cured should not be excluded from consideration for transplant. ISHLT guidelines have recommended stratifying each patient with pretransplant malignancy as to his or her risk of tumor recurrence and that cardiac transplantation should be considered when tumor recurrence is low based on tumor type, response to therapy, and negative metastatic workup. The guidelines also recommend that the specific amount of time to wait for transplant

after neoplasm remission will depend on these factors and no arbitrary time period for observation should be used.

### Human Immunodeficiency Virus Infection

Solid-organ transplant for patients who are HIV-positive has historically been controversial. The availability of highly active antiretroviral therapy has changed the natural history of the disease. Agüero et al. (2016) reported on a review of heart transplants among HIV-infected patients. (46) In this review, since 2001, 12 heart transplantations in HIV-infected patients were reported and 3 patients acquired HIV after heart transplantation. Fourteen (93%) of these 15 patients were younger than 50 years of age, with cluster of differentiation 4 (CD4) counts greater than 200 cells/mm<sup>3</sup>, and all recipients were taking antiretroviral therapy. Thirteen were alive with normal graft function at the end of follow-up. One patient had suboptimal adherence to antiretroviral therapy and died of multiorgan failure. The cause of death in the other patient was not reported. (47)

There are few data directly comparing outcomes for patients with and without HIV. In 2021, Doberne et al. compared survival outcomes of cardiac transplantation in HIV-positive recipients with HIV-negative recipients. (48) Utilizing UNOS data on first-time heart transplant recipients and their donors between January 2005 and June 2019, a total of 75 HIV-positive transplant recipients and 29,848 HIV-negative recipients were included in an analysis. Results revealed no difference in 30-day, 1-year, and 5-year survival of HIV-positive versus HIV-negative heart transplant recipients. However, HIV-positive recipients had significantly longer median lengths of hospital stays (18 vs. 15 days;  $p=.006$ ), rate of acute rejection during initial hospitalization (38.7% vs. 17.7%;  $p<.001$ ), and rate of anti-rejection treatment administration (26.7% vs. 10.4%;  $p<.001$ ).

Current OPTN policy permits HIV-positive transplant candidates. (49)

The British HIV Association and the British Transplantation Society (2017) updated their guidelines on kidney transplantation in patients with HIV disease. (50) These criteria may be extrapolated to other organs:

- Adherent with treatment, particularly antiretroviral therapy
- CD4 count greater than 100 cells/mL (ideally >200 cells/mL) for at least 3 months
- Undetectable HIV viremia (<50 HIV-1 RNA copies/mL) for at least 6 months
- No opportunistic infections for at least 6 months
- No history of progressive multifocal leukoencephalopathy, chronic intestinal cryptosporidiosis, or lymphoma.

### Age

The maximum acceptable age for heart transplantation is uncertain. While the maximum recipient age for heart transplantation had been set at 55 years, with more evidence of comparable survival rates among the older population following heart transplantation, transplant centers are accepting older recipients. Currently, the upper age limit for heart transplant candidates is generally defined by transplant centers.

Jamil et al. (2017) conducted a retrospective study of age as it relates to primary graft dysfunction after heart transplantation. (51) Of the 255 heart transplants studied, 70 (27%) recipients were 65 years and older and 185 were younger; there were no significant differences in posttransplant morbidity (all  $p > .12$ ) or 1-year survival between groups ( $p = .88$ ). The incidence of moderate or severe primary graft dysfunction was lower among the older patients (6%) than in the younger (16%;  $p = .037$ ). Study limitations included the single-center design, lack of data on long-term survival, and the potential for selection bias in retrospective studies.

Cooper et al. (2016) analyzed UNOS data to assess the long-term outcomes of older recipients of orthotopic heart transplantation (OHT) in the U.S. between 1987 and 2014. (52) During this period, 50,432 patients underwent OHT; 71.8% ( $n = 36,190$ ) were 18 to 59 years of age, 26.8% ( $n = 13,527$ ) were 60 to 69 years of age, and 1.4% ( $n = 715$ ) were 70 years of age or older. The 5-year mortality rate was 26.9% for recipients 18 to 59 years, 29.3% for recipients 60 to 69 years, and 30.8% for recipients 70 years of age and older. Survival between the oldest group and the 60 to 69-year-old group did not differ significantly ( $p = .48$ ).

Awad et al. (2016) reported on a single-center retrospective review of 704 adults who underwent heart transplantation from 1988 to 2012 to investigate the mortality and morbidity rates of heart transplantations among recipients 70 years of age and older ( $n = 45$ ) compared with recipients younger than 70 years of age ( $n = 659$ ). (53) The older and younger groups had similar 1-year (93.0% vs. 92.1%;  $p = .79$ ), 5-year (84.2% vs. 73.4%;  $p = .18$ ), and 10-year (51.2% vs. 50.2%;  $p = .43$ ) survival rates, respectively.

Kilic et al. (2012) analyzed UNOS data for 5,330 patients age 60 and older (mean age, 63.7 years) who underwent heart transplantation between 1995 and 2004. (54) A total of 3492 (65.5%) patients survived to 5 years. In multivariate analysis, statistically significant predictors of 5-year survival included younger age (OR, 0.97; 95% CI, 0.95 to 1.00), younger donor age (OR, 0.99; 95% CI, 0.99 to 1.00), White race (OR, 1.23; 95% CI, 1.02 to 1.49), shorter ischemic time (OR, 0.93; 95% CI, 0.87 to 0.99), and lower serum creatinine level (OR, 0.92; 95% CI, 0.87 to 0.98). In addition, hypertension, diabetes, and mechanical ventilation each significantly decreased the odds of surviving to 5 years. Patients with 2 or more of these factors had a 12% lower rate of 5-year survival than those with none.

### Pulmonary Hypertension

Findings from several studies have suggested that patients with pulmonary hypertension who successfully undergo treatment can subsequently have good outcomes after a heart transplant. (55-58) For example, Tsukashita et al. (2015) retrospectively compared the effect of continuous-flow LVAD support on pulmonary hypertension with posttransplantation outcomes among 227 potential OHT candidates with preexisting pulmonary hypertension. (59) Patients were divided into 2 groups based on preimplantation pulmonary vascular resistance (PVR): low ( $< 5$  Wood units) ( $n = 182$ ) and high ( $\geq 5$  Wood units) ( $n = 45$ ). After LVAD implantation, PVR in the high PVR group decreased significantly (7.13 Wood units to 2.82 Wood units,  $p < .001$ ) to a level similar to that seen in the low PVR group (2.70 Wood units,  $p = .91$ ) and remained low after

heart transplantation. The mean follow-up after OHT was 3.5 years (range, 1 month to 9.3 years). The in-hospital mortality rate after OHT was significantly higher in the high PVR group (20.7%) than in the low PVR group (5.8%;  $p<.05$ ). The survival rates at 3 years post-OHT were 85.0% for the low PVR group and 79.0% for the high PVR group ( $p=.45$ ).

De Santo et al. (2012) reported on 31 consecutive patients diagnosed with unresponsive pulmonary hypertension at baseline after right heart catheterization. (55) After 12 weeks of treatment with oral sildenafil, right heart catheterization showed reversibility of pulmonary hypertension, allowing patients to be listed for a heart transplant. Oral sildenafil treatment resumed following the transplant. One patient died in the hospital. A right heart catheterization at 3 months posttransplant showed normalization of the pulmonary hemodynamic profile, thereby allowing weaning from sildenafil in the 30 patients who survived hospitalization. The reversal of pulmonary hypertension was confirmed at 1 year in the 29 surviving patients. Similarly, in a study by Perez-Villa et al (2013), 22 patients considered high-risk for a heart transplant due to severe pulmonary hypertension were treated with bosentan. (56) After 4 months of treatment, the mean PVR decreased from 5.6 to 3.4 Wood units. In a similar group of 9 patients who refused participation and served as controls, mean PVR during this time increased from 4.6 to 5.5 Wood units. After bosentan therapy, 14 patients underwent heart transplantation, and the 1-year survival rate was 93%.

### Renal Insufficiency

A retrospective report by Arshad et al. (2019) compared renal outcomes and survival in patients who received an LVAD ( $n=45$ ) or heart transplant ( $n=58$ ). (60) The eGFR was similar between LVAD and transplant groups on day 30 after the procedure (75.1 mL/min/1.73 m<sup>2</sup> and 65.8 mL/min/1.73 m<sup>2</sup>, respectively;  $p=.057$ ), and significantly higher with LVAD versus transplant at 6 months (68.3 mL/min/1.73 m<sup>2</sup> and 59.4 mL/min/1.73 m<sup>2</sup>;  $p=.046$ ) and 1 year (68.3 mL/min/1.73 m<sup>2</sup> and 56.8 mL/min/1.73 m<sup>2</sup>;  $p=.15$ ). Survival rates were similar between LVAD and transplant groups at 1 year (84.4% and 81.0%, respectively;  $p=.540$ ) and 2 years (78.3% and 78.8%, respectively;  $p=.687$ ) after the procedure.

Another retrospective report by Kolsrud et al. (2018) investigated the association between post-heart transplantation and measured GFR as a risk factor for death and/or end-stage renal disease. (61) During the first year after heart transplant, 416 adults showed a 12% mean drop in measured GFR compared with preoperative values and long-term survival was significantly worse in patients who experienced a 25% or greater decrease in measured GFR during the first post-transplantation year (HR, 1.62; 95% CI, 1.04 to 2.53;  $p=.03$ ). Preoperative measured GFR was not predictive of mortality or end-stage renal disease, but older patients (HR, 1.03; 95% CI, 1.02 to 1.04;  $p<.001$ ) or patients with a ventricular assist device (HR, 2.23; 95% CI, 1.43 to 3.46;  $p<.001$ ) were predictors of death. The authors concluded that pretransplantation measured GFR was not predictive of mortality or end-stage renal disease after heart transplantation, but in this select patient population, a simultaneous or late-stage concomitant kidney transplant was necessary. Patients who experienced a 25% or greater measured eGFR decrease had the poorest prognosis. Study limitations included selection bias of patients, the retrospective study design, the exclusion of the sickest patients eligible undergoing post-heart transplantation,

changes in ventricular assist device and concomitant kidney transplant methods over time, and the small sample size studied.

The 2016 ISHLT criteria for heart transplantation recommended irreversible renal dysfunction ( $\text{eGFR} < 30 \text{ mL/min/1.73 m}^2$ ) as a relative contraindication for heart transplantation alone. The cutoff for eGFR in the previous recommendation was  $35 \text{ mL/min/1.73 m}^2$ . Hong et al. (2016) assessed 17,459 adult OHT recipients with results between 2001 and 2009 in the UNOS database to determine whether survival after OHT was associated with pretransplant eGFR and to define ranges of pretransplant eGFR associated with differences in posttransplant survival. (62) Posttransplant graft survival in the group with an eGFR less than  $34 \text{ mL/min/1.73 m}^2$  was significantly worse than in the groups with an eGFR of 35 to  $49 \text{ mL/min/1.73 m}^2$  or an eGFR greater than  $49 \text{ mL/min/1.73 m}^2$  ( $p < .001$ ). Median survival in the 3 groups was 8.2 years, 10.0 years, and 10.3 years, respectively. At 3 months, graft survival rates were 82.1%, 90.7%, and 94.0% in the groups with an eGFR less than  $34 \text{ mL/min/1.73 m}^2$ , an eGFR of 35 to  $49 \text{ mL/min/1.73 m}^2$ , and an eGFR greater than  $49 \text{ mL/min/1.73 m}^2$ , respectively. In multivariable logistic regression analysis, an eGFR less than  $34 \text{ mL/min/1.73 m}^2$  and an eGFR of 35 to  $49 \text{ mL/min/1.73 m}^2$  were significant risk factors for death at 1 year ( $p < .001$ ). Rossano et al (2016) also reported eGFR to be an independent risk factor for 1-, 5-, and 10-year posttransplant mortality among pediatric transplant recipients (described in the Pediatric Considerations section for survival after heart transplant). (30)

#### Children With Intellectual Disability

Considering the shortage of available donor organs, heart transplantation in children with intellectual disability has been debated. In 2016, ISHLT removed explicit mention of "mental retardation" as a relative contraindication to heart transplantation from its official guidelines. Multiple studies in recent years have examined whether intellectual disability in children is associated with significantly lower survival following heart transplantation compared with children without intellectual disability.

Goel et al. (2017) conducted a retrospective cohort study using UNOS data from 2008 to 2015 to evaluate the prevalence and outcomes of heart transplantation in this population. (63) Intellectual disability was assessed by using the cognitive development, academic progress, and academic level (5-point Likert scale scores for each of those) reported by transplant centers to UNOS. There were 565 pediatric ( $< 19$  years) patients with definite ( $n = 131$ ) or probable ( $n = 434$ ) intellectual disability who received their first heart transplant, accounting for 22.4% of all first pediatric heart transplants ( $n = 2524$ ). Intellectual disability was associated with a prolonged waitlist time ( $p < .001$ ). Patient survival rates at 1 and 3 years, respectively, were 88.9% and 86.0% for the definite intellectual disability group, 91.6% and 82.4% for the probable intellectual disability group, and 91.8% and 86.2% for the no intellectual disability group. Patient survival did not differ between groups at any time posttransplant ( $p = .578$ ). Intellectual disability status at listing was not associated with graft mortality hazards in univariate and multivariate analyses.

Wightman et al. (2017) performed a retrospective cohort analysis of 1204 children receiving a first isolated heart transplant for whom cognitive and educational data were available in the UNOS dataset between 2008 and 2013. (64) Children were categorized as "definitely cognitive delay/impairment" by their transplant center using the Likert scales for cognitive development. All other recipients were classified as "no intellectual disability." Kaplan-Meier curves and log-rank tests did not suggest a significant difference in graft survival during the first 4 years after transplantation ( $p=.07$ ), however, they did suggest poorer patient survival among the intellectual disability group during the first 4 years following transplantation ( $p=.05$ ). In an unadjusted Cox regression, intellectual disability was associated with poorer graft (HR, 1.66; 95% CI, 1.01 to 2.72;  $p=.05$ ) and patient survival (HR, 1.71; 95% CI, 0.99 to 2.94;  $p=.05$ ). However, after adjusting for covariates, there was no association between intellectual disability and graft survival (HR, 0.95; 95% CI, 0.49 to 1.88;  $p=.89$ ) or patient survival (HR, 0.80; 95% CI, 0.36 to 1.75;  $p=.58$ ). Wightman et al. (2021) also investigated the prevalence and long-term outcomes of initial kidney, liver, and heart transplants from 2008 to 2017 using UNOS data in children with an intellectual disability. (65) During this study period, children with definite intellectual disability accounted for 324 (9%) of 3722 initial heart transplant recipients. In these patients, intellectual disability was not significantly associated with patient or graft survival.

Prendergast et al. (2017) assessed the impact of cognitive delay on pediatric heart transplantation outcomes using academic progress as a surrogate for cognitive performance among pediatric heart transplant recipients (2004 to 2014) with data reporting academic progress in the OPTN database ( $n=2245$ ). (66) Of the patients with complete academic progress data, 1707 (76%) were within 1 grade level of peers, 269 (12%) had delayed grade level, and 269 (12%) required special education. There was no significant difference in posttransplant survival between patients within 1 grade level of peers and those who required special education. However, patients with delayed grade level demonstrated worse post-transplant survival than patients within 1 grade level of peers and those who required special education ( $p<.001$ ). Delayed grade level remained as an independent predictor of posttransplant graft loss (adjusted HR, 1.4; 95% CI, 1.02 to 1.79;  $p=.03$ ) in multivariate analysis. The authors conducted a secondary analysis substituting cognitive delay for academic progress; patients were divided into 2 groups based on whether any concerns for a cognitive delay (questionable, probable, or definite) were ever reported at the time of heart transplantation or during follow-up (1176 with cognitive delay, 1783 with no documented cognitive delay). There was no significant difference in posttransplant graft survival based on the presence of cognitive delay ( $p=.57$ ). Cognitive delay remained a statistically nonsignificant predictor in multivariate analysis (adjusted HR, 1.01; 95% CI, 0.83 to 1.22;  $p=.953$ ).

Because these studies assessed patients who received transplants and did not evaluate children who were refused listing by a transplant center or never referred to a transplant center, the prevalence of intellectual disability among potential candidates of heart transplantation might have been underestimated. With low-risk intellectual disability patients receiving heart transplant and individuals with intellectual disability and other high-risk conditions being excluded, results might also have a positive selection bias.

## Summary of Evidence

For individuals who have end-stage heart failure who receive a heart transplant, the evidence includes retrospective studies and registry data. Relevant outcomes are overall survival (OS), symptoms, and morbid events. Heart transplant remains a viable treatment for those with severe heart dysfunction despite appropriate medical management with medication, surgery, or medical devices. Given the exceedingly poor survival rates without transplantation for these patients, evidence of post-transplant survival is sufficient to demonstrate that heart transplantation provides a survival benefit. Heart transplantation is contraindicated for patients for whom the procedure is expected to be futile due to comorbid disease or for whom post-transplantation care is expected to worsen comorbid conditions significantly. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have had a prior heart transplant complicated by graft failure or severe dysfunction of the heart who receive a heart retransplant, the evidence includes systematic reviews, retrospective studies, and registry data. Relevant outcomes are OS, symptoms, and morbid events. Despite improvements in the prognosis for many patients with graft failure, cardiac allograft vasculopathy, and severe dysfunction of the transplanted heart, heart retransplant remains a viable treatment for those whose severe symptoms persist despite treatment with other medical or surgical remedies. Given the exceedingly poor survival rates without retransplantation for patients who have exhausted other treatments, evidence of post-transplant survival is sufficient to demonstrate that heart retransplantation provides a survival benefit in appropriately selected patients. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

## Practice Guidelines and Position Statements

### American College of Cardiology Foundation et al.

Heart failure guidelines from the American College of Cardiology Foundation, the American Heart Association, and the Heart Failure Society of America were updated in 2022. (11)

Recommendations for cardiac transplantation by the joint committee were as follows:

- "For selected patients with advanced HF [heart failure] despite GDMT [guideline-directed medical therapy], cardiac transplantation is indicated to improve survival and QOL [quality of life] (class of recommendation, 1 [strong]; level of evidence, C-LD [limited data, including registry or observational studies])."
- "In patients with stage D (advanced) HF despite GDMT, cardiac transplantation provides intermediate economic value (value statement: intermediate value)."

### American Heart Association

In 2007, the American Heart Association indicated that, based on level B (nonrandomized studies) or level C (consensus opinion of experts) evidence, heart transplantation is indicated for pediatric patients as therapy for the following indications: (67)

- Stage D heart failure (interpreted as abnormal cardiac structure and/or function, continuous infusion of intravenous inotropes, or prostaglandin E<sub>1</sub> to maintain patency of a

ductus arteriosus, mechanical ventilatory and/or mechanical circulatory support) associated with systemic ventricular dysfunction in patients with cardiomyopathies or previous repaired or palliated congenital heart disease,

- Stage C heart failure (interpreted as abnormal cardiac structure and/or function and past or present symptoms of heart failure) associated with pediatric heart disease and severe limitation of exercise and activity, in patients with cardiomyopathies or previously repaired or palliated congenital heart disease and heart failure associated with significant growth failure attributed to heart disease, pediatric heart disease with associated near sudden death and/or life-threatening arrhythmias untreatable with medications or an implantable defibrillator, or in pediatric restrictive cardiomyopathy disease associated with reactive pulmonary hypertension;

The guideline states that heart transplantation is feasible in the presence of other indications for heart transplantation, "in patients with pediatric heart disease and an elevated pulmonary vascular resistance index  $>6$  Woods units/ $m^2$  and/or a transpulmonary pressure gradient  $>15$  mm Hg if administration of inotropic support or pulmonary vasodilators can decrease pulmonary vascular resistance to  $<6$  Woods units/ $m^2$  or the transpulmonary gradient to  $<15$  mm Hg."

#### International Society for Heart and Lung Transplantation

In 2004, the International Society for Heart and Lung Transplantation (ISHLT) recommended that children with the following conditions be evaluated for heart transplantation (Table 3). (68)

**Table 3. Recommendations for Pediatric Heart Transplant**

| Recommendation                                                                                                                                                                                                                       | LOE |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Diastolic dysfunction that is refractory to optimal medical/surgical management because they are at high risk of developing pulmonary hypertension and of sudden death                                                               | B   |
| Advanced systemic right ventricular failure (Heart Failure Stage C described as patients with underlying structural or functional heart disease and past or current symptoms of heart failure) that is refractory to medical therapy | C   |

LOE B is based on a single randomized trial or multiple nonrandomized trials; LOE C is based primarily on expert consensus opinion.

LOE: level of evidence.

In 2024, the ISHLT published an update to guidelines for the evaluation and care of cardiac transplant candidates that replaces previous versions from 2016 and 2006. (69) Select relevant recommendations regarding indications for heart transplantation and retransplantation are listed in Table 4.

**Table 4. Recommendations for Heart Transplantation and Retransplantation**

| Recommendation | COR <sup>a</sup> | LOE <sup>b</sup> |
|----------------|------------------|------------------|
|----------------|------------------|------------------|

|                                                                                                                                                                                                                                                                                                                                     |    |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|------|
| In patients with HF, when consistent with the patient's goals of care, the presence of clinical indicators of AdvHF should trigger evaluation for AdvHF therapies, including HT.                                                                                                                                                    | 1  | B-NR |
| In patients with AdvHF aged $\leq 70$ years, evaluation for HT is recommended.                                                                                                                                                                                                                                                      | 1  | B-NR |
| In patients with AdvHF who are over 70 years of age, evaluation for HT may be considered in carefully selected patients depending on their functional status and control of comorbidities, taking into account the need for specialized post-transplant care for older patients, including the need for tailored immunosuppression. | 2b | B-NR |
| In heart transplant recipients with ISHLT Grade 3 CAV, evaluation for retransplantation is reasonable.                                                                                                                                                                                                                              | 2a | B-NR |
| In heart transplant recipients with the following, the benefit of retransplantation is not well established: a. Graft failure due to active rejection; b. Advanced age; c. Need for DMCS as a bridge to retransplantation.                                                                                                          | 2b | C-LD |

AdvHF: advanced HF; CAV: cardiac allograft vasculopathy; COR: class of recommendation; DMCS: durable mechanical circulatory support; HF: heart failure; HT: heart transplantation; ISHLT: International Society for Heart and Lung Transplantation; LOE: level of evidence; RCT: randomized controlled trial.

<sup>a</sup> Class (strength) of recommendation - Class 1 (strong): Benefit >>> Risk; Class 2a (moderate): Benefit >> Risk; Class 2b (weak): Benefit > Risk; Class 3: no benefit (moderate): Benefit = Risk; Class 3: harm (strong): Risk > Benefit.

<sup>b</sup> Level (quality) of evidence - Level A: High-quality evidence from more than 1 RCT OR meta-analyses of high quality RCTs OR 1 or more RCT corroborated by high-quality registry studies; Level B-R (Randomized): Moderate-quality evidence from 1 or more RCTs OR meta-analyses of moderate-quality RCTs; Level B-NR (Nonrandomized): Moderate-quality evidence from 1 or more well-designed, well-executed nonrandomized studies, observational studies, or registry studies OR meta-analyses of such studies; Level C-LD (Limited Data): Randomized or nonrandomized observational or registry studies with limitations of design or execution OR meta-analyses of such studies OR physiological or mechanistic studies in human subjects; Level C-EO (Expert Opinion): Consensus of expert opinion based on clinical experience.

In 2023, the guidelines from ISHLT on the care of heart transplant recipients were updated from the previous 2010 version. (70) Recommendations made on cardiac retransplantation are below:

- "Retransplantation is indicated in children with at least moderate systolic heart allograft dysfunction and/or severe diastolic dysfunction and at least moderate CAV (cardiac allograft vasculopathy; Class I, level of evidence, B)."
- "Evaluation for retransplantation should be considered in adults and children with severe CAV and allograft dysfunction in the absence of contraindications for repeat HF (Class IIa, level of evidence, B)."
- "It is reasonable to consider retransplantation in children with normal heart allograft function and severe CAV (Class IIa, level of evidence B)."

## Medicare National Coverage

Cardiac transplantation is covered under Medicare when performed in a facility approved by Medicare. (71) The Centers for Medicare & Medicaid Services has stated that, under certain limited cases, exceptions to the criteria may be warranted if there is justification and if the facility ensures safety and efficacy objectives.

### Ongoing and Unpublished Clinical Trials

A search of ClinicalTrials.gov in July 2025 did not identify any ongoing or unpublished trials that would likely influence this policy.

## Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member's benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

|                    |                     |
|--------------------|---------------------|
| <b>CPT Codes</b>   | 33940, 33944, 33945 |
| <b>HCPSC Codes</b> | S2152               |

\*Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.

## References

1. Black CK, Termanini KM, Aguirre O, et al. Solid organ transplantation in the 21st century. *Ann Transl Med.* Oct 2018; 6(20):409. PMID 30498736
2. United Network for Organ Sharing (UNOS). Data and trends (2025). Available at <<https://www.unos.org>> (accessed July 30, 2025).
3. Organ Procurement and Transplantation Network (OPTN). National data. 2025. Available at <<https://www.optn.transplant.hrsa.gov>> (accessed July 30, 2025).
4. Rose SW, Strackman BW, Gilbert ON, et al. Disparities by Sex, Race, and Ethnicity in Use of Left Ventricular Assist Devices and Heart Transplants Among Patients With Heart Failure With Reduced Ejection Fraction. *J Am Heart Assoc.* Jan 16 2024; 13(2):e031021. PMID 38166429
5. Lietz K, Miller LW. Improved survival of patients with end-stage heart failure listed for heart transplantation: analysis of organ procurement and transplantation network/U.S. United Network of Organ Sharing data, 1990 to 2005. *J Am Coll Cardiol.* Sep 25 2007; 50(13):1282-1290. PMID 17888847
6. Johnson MR, Meyer KH, Haft J, et al. Heart transplantation in the United States, 1999-2008. *Am J Transplant.* Apr 2010; 10(4 Pt 2):1035-1046. PMID 20420651
7. Bakhtiyar SS, Godfrey EL, Ahmed S, et al. Survival on the heart transplant waiting list. *JAMA Cardiol.* Nov 01 2020; 5(11):1227-1235. PMID 32785619

8. Alshawabkeh L, Opatowsky AR, Carter KD, et al. Disparities in wait-list outcomes for adults with congenital heart disease listed for heart transplantation before and since revision of status I listing. *Am J Cardiol*. Nov 15 2018; 122(10):1761-1764. PMID 30236623
9. Alshawabkeh LI, Hu N, Carter KD, et al. Wait-list outcomes for adults with congenital heart disease listed for heart transplantation in the U.S. *J Am Coll Cardiol*. Aug 30 2016; 68(9):908-917. PMID 27561764
10. Magnetta DA, Godown J, West S, et al. Impact of the 2016 revision of US Pediatric Heart Allocation Policy on waitlist characteristics and outcomes. *Am J Transplant*. Dec 2019; 19(12):3276-3283. PMID 31544351
11. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. May 03 2022; 79(17):e263-e421. PMID 35379503
12. Aaronson KD, Schwartz JS, Chen TM, et al. Development and prospective validation of a clinical index to predict survival in ambulatory patients referred for cardiac transplant evaluation. *Circulation*. Jun 17 1997; 95(12):2660-2667. PMID 9193435
13. Alla F, Briancon S, Juilliere Y, et al. Differential clinical prognostic classifications in dilated and ischemic advanced heart failure: the EPICAL study. *Am Heart J*. 2000; 139(5):895-904. PMID 10783225
14. Hansen A, Haass M, Zugck C, et al. Prognostic value of Doppler echocardiographic mitral inflow patterns: implications for risk stratification in patients with chronic congestive heart failure. *J Am Coll Cardiol*. 2001; 37(4):1049-1055. PMID 11263607
15. Lee DS, Austin PC, Rouleau JL, et al. Predicting mortality among patients hospitalized for heart failure: derivation and validation of a clinical model. *JAMA*. 2003; 290(19):2581-2587. PMID 14625335
16. Levy WC, Mozaffarian D, Linker DT, et al. The Seattle Heart Failure Model: prediction of survival in heart failure. *Circulation*. Mar 21 2006; 113(11):1424-1433. PMID 16534009
17. Gorodeski E, Chu EC, Chow CH, et al. Application of the Seattle Heart Failure Model in Ambulatory Patients Presented to an Advanced Heart Failure Therapeutics Committee. *Circ Heart Fail*. Nov 2010; 3(6):706-714. PMID 20798278
18. Ketchum ES, Moorman AJ, Fishbein DP, et al. Predictive value of the Seattle Heart Failure Model in patients undergoing left ventricular assist device placement. *J Heart Lung Transplant*. Sep 2010; 29(9):1021-1025. PMID 20558086
19. Nutter AL, Tanawuttiwat T, Silver MA. Evaluation of 6 prognostic models used to calculate mortality rates in elderly heart failure patients with a fatal heart failure admission. *Congest Heart Fail*. Sep-Oct 2010; 16(5):196-201. PMID 20887615
20. Kalogeropoulos AP, Georgiopoulos VV, Giamouzis G, et al. Utility of the Seattle Heart Failure Model in patients with advanced heart failure. *J Am Coll Cardiol*. Jan 27 2009; 53(4):334-342. PMID 19161882
21. May HT, Horne BD, Levy WC, et al. Validation of the Seattle Heart Failure Model in a community-based heart failure population and enhancement by adding B-type natriuretic peptide. *Am J Cardiol*. Aug 15 2007; 100(4):697-700. PMID 17697831

22. Virani SS, Alonso A, Aparicio HJ, et al. Heart Disease and Stroke Statistics-2021 Update: A Report From the American Heart Association. *Circulation*. Feb 23 2021; 143(8):e254-e743. PMID 33501848
23. Lund LH, Edwards LB, Dipchand AI, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirty-third Adult Heart Transplantation Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant*. Oct 2016; 35(10):1158-1169. PMID 27772668
24. Jaramillo N, Segovia J, Gomez-Bueno M, et al. Characteristics of patients with survival longer than 20 years following heart transplantation. *Rev Esp Cardiol*. Oct 2013;66(10):797-802. PMID 23932221
25. Nguyen VP, Mahr C, Mokadam NA, et al. The benefit of donor-recipient matching for patients undergoing heart transplantation. *J Am Coll Cardiol*. Apr 04 2017; 69(13):1707-1714. PMID 28359517
26. Rana A, Gruessner A, Agopian VG, et al. Survival benefit of solid-organ transplant in the United States. *JAMA Surg*. Mar 1 2015; 150(3):252-259. PMID 25629390
27. Kilic A, Weiss ES, George TJ, et al. What predicts long-term survival after heart transplantation? An analysis of 9,400 ten-year survivors. *Ann Thorac Surg*. Mar 2012; 93(3):699-704. PMID 22226494
28. Dipchand AI, Kirk R, Mahle WT, et al. Ten yr of pediatric heart transplantation: a report from the Pediatric Heart Transplant Study. *Pediatr Transplant*. Mar 2013; 17(2):99-111. PMID 23442098
29. Auerbach SR, Richmond ME, Chen JM, et al. Multiple risk factors before pediatric cardiac transplantation are associated with increased graft loss. *Pediatr Cardiol*. Jan 2012; 33(1):49-54. PMID 21892650
30. Rossano JW, Dipchand AI, Edwards LB, et al. The Registry of the International Society for Heart and Lung Transplantation: Nineteenth Pediatric Heart Transplantation Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant*. Oct 2016; 35(10):1185-1195. PMID 27772670
31. Savla J, Lin KY, Lefkowitz DS, et al. Adolescent age and heart transplantation outcomes in myocarditis or congenital heart disease. *J Heart Lung Transplant*. Sep 2014; 33(9):943-949. PMID 24929645
32. Almond CSD, Thiagarajan RR, Piercey GE, et al. Waiting list mortality among children listed for heart transplantation in the United States. *Circulation*. Feb 10 2009; 119(5):717-727. PMID 19171850
33. Tjang YS, Tenderich G, Hornik L, et al. Cardiac retransplantation in adults: an evidence-based systematic review. *Thorac Cardiovasc Surg*. Sep 2008; 56(6):323-327. PMID 18704853
34. Zhu Y, Shudo Y, Lingala B, et al. Outcomes after heart retransplantation: A 50-year single-center experience. *J Thorac Cardiovasc Surg*. Feb 2022; 163(2):712-720.e6. PMID 32798029
35. Saito A, Novick RJ, Kiaii B, et al. Early and late outcomes after cardiac retransplantation. *Can J Surg*. Feb 2013; 56(1):21-26. PMID 23187039
36. Miller RJH, Clarke BA, Howlett JG, et al. Outcomes in patients undergoing cardiac retransplantation: A propensity matched cohort analysis of the UNOS Registry. *J Heart Lung Transplant*. Oct 2019; 38(10):1067-1074. PMID 31378576

37. Goldraich LA, Stehlik J, Kucheryavaya AY, et al. Retransplant and medical therapy for cardiac allograft vasculopathy: international society for heart and lung transplantation registry analysis. *Am J Transplant*. Jan 2016; 16(1):301-309. PMID 26274617
38. Belli E, Leoni Moreno JC, Hosenpud J, et al. Preoperative risk factors predict survival following cardiac retransplantation: analysis of the United Network for Organ Sharing database. *J Thorac Cardiovasc Surg*. Jun 2014; 147(6):1972-1977, 197.e1971. PMID 24636155
39. Friedland-Little JM, Gajarski RJ, Yu S, et al. Outcomes of third heart transplants in pediatric and young adult patients: analysis of the United Network for Organ Sharing database. *J Heart Lung Transplant*. Sep 2014; 33(9):917-923. PMID 24861821
40. Bock MJ, Nguyen K, Malerba S, et al. Pediatric cardiac retransplantation: Waitlist mortality stratified by age and era. *J Heart Lung Transplant*. Apr 2015; 34(4):530-537. PMID 25016920
41. Conway J, Manlhiot C, Kirk R, et al. Mortality and morbidity after retransplantation after primary heart transplant in childhood: an analysis from the registry of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. Mar 2014; 33(3):241-251. PMID 24462559
42. Mistiaen WP. Heart transplantation in patients with previous malignancy. An overview. *Acta Cardiol*. Apr 2015; 70(2):123-130. PMID 26148371
43. Yoosabai A, Mehta A, Kang W, et al. Pretransplant malignancy as a risk factor for posttransplant malignancy after heart transplantation. *Transplantation*. Feb 2015; 99(2):345-350. PMID 25606783
44. Oliveira GH, Hardaway BW, Kucheryavaya AY, et al. Characteristics and survival of patients with chemotherapy- induced cardiomyopathy undergoing heart transplantation. *J Heart Lung Transplant*. Aug 2012; 31(8):805-810. PMID 22551930
45. Sigurdardottir V, Bjortuft O, Eiskjaer H, et al. Long-term follow-up of lung and heart transplant recipients with pre- transplant malignancies. *J Heart Lung Transplant*. Dec 2012; 31(12):1276-1280. PMID 23089300
46. Agüero F, Castel MA, Cocchi S, et al. An update on heart transplantation in human immunodeficiency virus- infected patients. *Am J Transplant*. Jan 2016; 16(1):21-28. PMID 26523614
47. Uriel N, Jorde UP, Cotarlan V, et al. Heart transplantation in human immunodeficiency virus- positive patients. *J Heart Lung Transplant*. Jul 2009; 28(7):667-669. PMID 19560693
48. Doberne JW, Jawitz OK, Raman V, et al. Heart Transplantation Survival Outcomes of HIV Positive and Negative Recipients. *Ann Thorac Surg*. May 2021; 111(5):1465-1471. PMID 32946847
49. Organ Procurement and Transplantation Network (OPTN). Organ Procurement and Transplantation Network Policies. 2025. Available at <<https://www.optn.transplant.hrsa.gov>> (accessed December 8, 2025).
50. Working Party of the British Transplantation Society. Kidney and Pancreas Transplantation in Patients with HIV. Second Edition (Revised). British Transplantation Society Guidelines. Macclesfield, UK: British Transplantation Society; 2017.
51. Jamil A, Qin H, Felius J, et al. Comparison of clinical characteristics, complications, and outcomes in recipients having heart transplants <65 years of age versus 65 years of age. *Am J Cardiol*. Sep 15 2017; 120(12):2207-2212. PMID 29056228

52. Cooper LB, Lu D, Mentz RJ, et al. Cardiac transplantation for older patients: Characteristics and outcomes in the septuagenarian population. *J Heart Lung Transplant*. Mar 2016; 35(3):362-369. PMID 26632028
53. Awad M, Czer LS, Mirocha J, et al. Similar mortality and morbidity of orthotopic heart transplantation for patients 70 years of age and older compared with younger patients. *Transplant Proc*. Oct 2016; 48(8):2782-2791. PMID 27788818
54. Kilic A, Weiss ES, Yuh DD, et al. Factors associated with 5-year survival in older heart transplant recipients. *J Thorac Cardiovasc Surg*. Feb 2012; 143(2):468-474. PMID 22248684
55. De Santo LS, Romano G, Maiello C, et al. Pulmonary artery hypertension in heart transplant recipients: how much is too much? *Eur J Cardiothorac Surg*. Nov 2012; 42(5):864-869; discussion 869-870. PMID 22402452
56. Perez-Villa F, Farrero M, Cardona M, et al. Bosentan in heart transplantation candidates with severe pulmonary hypertension: efficacy, safety and outcome after transplantation. *Clin Transplant*. Jan-Feb 2013; 27(1):25-31. PMID 22861120
57. Pons J, Leblanc MH, Bernier M, et al. Effects of chronic sildenafil use on pulmonary hemodynamics and clinical outcomes in heart transplantation. *J Heart Lung Transplant*. Dec 2012; 31(12):1281-1287. PMID 23127754
58. Bedanova H, Orban M, Vrsansky D, et al. Impact of pulmonary hypertension on early hemodynamics, morbidity and mortality after orthotopic heart transplantation. A single center study. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*. Mar 2013; 157(1):35-40. PMID 23073529
59. Tsukashita M, Takayama H, Takeda K, et al. Effect of pulmonary vascular resistance before left ventricular assist device implantation on short- and long-term post-transplant survival. *J Thorac Cardiovasc Surg*. Nov 2015; 150(5):1352-1360, 1361.e1-2. PMID 26253875
60. Arshad A, Kew EP, Lim S. Comparison of renal outcomes in patients with left ventricular assist device and heart transplantation. *Transplant Proc*. Dec 2019; 51(10):3395-3398. PMID 31810507
61. Kolsrud O, Karason K, Holmberg E, et al. Renal function and outcome after heart transplantation. *J Thorac Cardiovasc Surg*. Dec 14 2018; 155(4):1593-1604.e1591. PMID 29338859
62. Hong KN, Merlo A, Chauhan D, et al. Evidence supports severe renal insufficiency as a relative contraindication to heart transplantation. *J Heart Lung Transplant*. Jul 2016; 35(7):893-900. PMID 27105687
63. Goel AN, Iyengar A, Schowengerdt K, et al. Heart transplantation in children with intellectual disability: An analysis of the UNOS database. *Pediatr Transplant*. Mar 2017; 21(2). PMID 27933693
64. Wightman A, Bartlett HL, Zhao Q, et al. Prevalence and outcomes of heart transplantation in children with intellectual disability. *Pediatr Transplant*. Mar 2017; 21(2). PMID 27801533
65. Wightman A, Bradford MC, Hsu E, et al. Prevalence and long-term outcomes of solid organ transplant in children with intellectual disability. *J Pediatr*. Aug 2021; 235:10-17.e4. PMID 33794218
66. Prendergast C, McKane M, Dodd DA, et al. The impact of cognitive delay on pediatric heart transplant outcomes. *Pediatr Transplant*. Mar 2017; 21(2). PMID 28191755

67. Canter CE, Shaddy RE, Bernstein D, et al. Indications for heart transplantation in pediatric heart disease: a scientific statement from the American Heart Association Council on Cardiovascular Disease in the Young; the Councils on Clinical Cardiology, Cardiovascular Nursing, and Cardiovascular Surgery and Anesthesia; and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation*. Feb 6 2007; 115(5):658- 676. PMID 17261651
68. Rosenthal D, Chrisant MR, Edens E, et al. International Society for Heart and Lung Transplantation: Practice guidelines for management of heart failure in children. *J Heart Lung Transplant*. Dec 2004; 23(12):1313-1333. PMID 15607659
69. Peled Y, Ducharme A, Kittleson M, et al. International Society for Heart and Lung Transplantation Guidelines for the Evaluation and Care of Cardiac Transplant Candidates- 2024. *J Heart Lung Transplant*. Oct 2024; 43(10):1529-1628.e54. PMID 39115488
70. Velleca A, Shullo MA, Dhital K, et al. The International Society for Heart and Lung Transplantation (ISHLT) guidelines for the care of heart transplant recipients. *J Heart Lung Transplant*. May 2023; 42(5):e1-e141. PMID 37080658
71. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for HEART TRANSPLANTS (260.9). 2008. Available at <<https://www.cms.gov>> (accessed July 30, 2025).

## Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been changed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

### Policy History/Revision

| Date       | Description of Change                                                                                                                                      |
|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 01/01/2026 | Document updated. Coverage unchanged. Added references 69 and 70; other(s) updated.                                                                        |
| 10/15/2024 | Document updated with literature review. Coverage unchanged. Added reference 4; others updated and some removed.                                           |
| 02/01/2024 | Document updated with literature review. Coverage unchanged. Added references 34, 67; others updated.                                                      |
| 12/01/2022 | Reviewed. No changes.                                                                                                                                      |
| 01/01/2022 | Document updated with literature review. Coverage unchanged. The following references were added/updated: 1, 2, 3, 6-9, 22, 24, 35, 47, 48, 59, 64 and 72. |

|            |                                                                                                                                                                                                                                                                                                                                                                                     |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 01/15/2021 | Reviewed. No changes.                                                                                                                                                                                                                                                                                                                                                               |
| 06/01/2020 | Document updated with literature review. The follow change was made to the Coverage section: The medically necessary statement was changed from including 1A, 1B, or 2 Status to now including the United Network for Organ Sharing (UNOS) guidelines for 1-6 Status. References 42, 51 and 56 were added and some references removed.                                              |
| 11/15/2018 | Reviewed. No changes.                                                                                                                                                                                                                                                                                                                                                               |
| 12/15/2017 | Document updated with literature review. The following change was made to Coverage: Updated terminology from "Status 7" to "Inactive Status".                                                                                                                                                                                                                                       |
| 03/01/2016 | Reviewed. No changes.                                                                                                                                                                                                                                                                                                                                                               |
| 06/01/2015 | Document updated with literature review. Coverage unchanged.                                                                                                                                                                                                                                                                                                                        |
| 12/01/2014 | Document updated with literature review. The following statements were added to coverage: 1) Heart retransplantation after a failed primary heart transplant may be considered medically necessary in patients who meet criteria for heart transplantation. 2) Heart transplantation is considered experimental, investigational and/or unproven in all other situations.           |
| 11/01/2013 | Document updated with literature review. Coverage changed to: Human heart transplant may be considered medically necessary in carefully selected patients with irreversible, refractory, and symptomatic end-stage heart failure who meet the United Network for Organ Sharing (UNOS) guidelines for 1A, 1B, or 2 Status and are not currently Status 7. CPT/HCPCS code(s) updated. |
| 04/15/2009 | Editorial revision to clarify end-stage cardiomyopathy coverage criteria; references revised                                                                                                                                                                                                                                                                                        |
| 06/01/2008 | Revised/updated entire document; this policy is no longer scheduled for routine literature and update                                                                                                                                                                                                                                                                               |
| 02/01/2005 | Revised/updated entire document                                                                                                                                                                                                                                                                                                                                                     |
| 09/01/1998 | Revised/updated entire document                                                                                                                                                                                                                                                                                                                                                     |
| 05/01/1996 | Medical policy number changed                                                                                                                                                                                                                                                                                                                                                       |
| 04/01/1996 | Revised/updated entire document                                                                                                                                                                                                                                                                                                                                                     |
| 01/01/1992 | Revised/updated entire document                                                                                                                                                                                                                                                                                                                                                     |
| 05/01/1990 | New medical document                                                                                                                                                                                                                                                                                                                                                                |